

Part IB — Methods

Theorems

Based on lectures by D. B. Skinner

Notes taken by Dexter Chua

Michaelmas 2015

These notes are not endorsed by the lecturers, and I have modified them (often significantly) after lectures. They are nowhere near accurate representations of what was actually lectured, and in particular, all errors are almost surely mine.

Self-adjoint ODEs

Periodic functions. Fourier series: definition and simple properties; Parseval's theorem. Equations of second order. Self-adjoint differential operators. The Sturm-Liouville equation; eigenfunctions and eigenvalues; reality of eigenvalues and orthogonality of eigenfunctions; eigenfunction expansions (Fourier series as prototype), approximation in mean square, statement of completeness. [5]

PDEs on bounded domains: separation of variables

Physical basis of Laplace's equation, the wave equation and the diffusion equation. General method of separation of variables in Cartesian, cylindrical and spherical coordinates. Legendre's equation: derivation, solutions including explicit forms of P_0 , P_1 and P_2 , orthogonality. Bessel's equation of integer order as an example of a self-adjoint eigenvalue problem with non-trivial weight.

Examples including potentials on rectangular and circular domains and on a spherical domain (axisymmetric case only), waves on a finite string and heat flow down a semi-infinite rod. [5]

Inhomogeneous ODEs: Green's functions

Properties of the Dirac delta function. Initial value problems and forced problems with two fixed end points; solution using Green's functions. Eigenfunction expansions of the delta function and Green's functions. [4]

Fourier transforms

Fourier transforms: definition and simple properties; inversion and convolution theorems. The discrete Fourier transform. Examples of application to linear systems. Relationship of transfer function to Green's function for initial value problems. [4]

PDEs on unbounded domains

Classification of PDEs in two independent variables. Well posedness. Solution by the method of characteristics. Green's functions for PDEs in 1, 2 and 3 independent variables; fundamental solutions of the wave equation, Laplace's equation and the diffusion equation. The method of images. Application to the forced wave equation, Poisson's equation and forced diffusion equation. Transient solutions of diffusion problems: the error function. [6]

Contents

0	Introduction	3
1	Vector spaces	4
2	Fourier series	5
2.1	Fourier series	5
2.2	Convergence of Fourier series	5
2.3	Differentiability and Fourier series	5
3	Sturm-Liouville Theory	6
3.1	Sturm-Liouville operators	6
3.2	Least squares approximation	6
4	Partial differential equations	7
4.1	Laplace's equation	7
4.2	Laplace's equation in the unit disk in \mathbb{R}^2	7
4.3	Separation of variables	7
4.4	Laplace's equation in spherical polar coordinates	7
4.4.1	Laplace's equation in spherical polar coordinates	7
4.4.2	Legendre Polynomials	7
4.4.3	Solution to radial part	7
4.5	Multipole expansions for Laplace's equation	7
4.6	Laplace's equation in cylindrical coordinates	7
4.7	The heat equation	7
4.8	The wave equation	7
5	Distributions	8
5.1	Distributions	8
5.2	Green's functions	8
5.3	Green's functions for initial value problems	8
6	Fourier transforms	9
6.1	The Fourier transform	9
6.2	The Fourier inversion theorem	9
6.3	Parseval's theorem for Fourier transforms	9
6.4	A word of warning	9
6.5	Fourier transformation of distributions	9
6.6	Linear systems and response functions	9
6.7	General form of transfer function	9
6.8	The discrete Fourier transform	9
6.9	The fast Fourier transform*	9
7	More partial differential equations	10
7.1	Well-posedness	10
7.2	Method of characteristics	10
7.3	Characteristics for 2nd order partial differential equations	10
7.4	Green's functions for PDEs on \mathbb{R}^n	10
7.5	Poisson's equation	10

0 Introduction

1 Vector spaces

2 Fourier series

2.1 Fourier series

2.2 Convergence of Fourier series

2.3 Differentiability and Fourier series

Theorem (Parseval's theorem).

$$(f, f) = \int_{-\pi}^{\pi} |f(\theta)|^2 d\theta = 2\pi \sum_{n \in \mathbb{Z}} |\hat{f}_n|^2$$

3 Sturm-Liouville Theory

3.1 Sturm-Liouville operators

Proposition. The eigenvalues of a Sturm-Liouville operator are real.

Proposition. Eigenfunctions with different eigenvalues (but same weight) are orthogonal.

Theorem. On a *compact* domain, the eigenvalues $\lambda_1, \lambda_2, \dots$ form a countably infinite sequence and are discrete.

Theorem. The eigenfunctions are complete: any function $f : [a, b] \rightarrow \mathbb{C}$ (obeying appropriate boundary conditions) can be expanded as

$$f(x) = \sum_n \hat{f}_n y_n(x),$$

where

$$\hat{f}_n = \int y_n^*(x) f(x) w(x) dx.$$

Theorem (Parseval's theorem II).

$$(f, f)_w = \sum_{n \in \mathbb{Z}} |\hat{f}_n|^2.$$

3.2 Least squares approximation

4 Partial differential equations

4.1 Laplace's equation

Proposition. Let Ω be a compact domain, and $\partial\Omega$ be its boundary. Let $f : \partial\Omega \rightarrow \mathbb{R}$. Then there is a unique solution to $\nabla^2\phi = 0$ on Ω with $\phi|_{\partial\Omega} = f$.

4.2 Laplace's equation in the unit disk in \mathbb{R}^2

4.3 Separation of variables

4.4 Laplace's equation in spherical polar coordinates

4.4.1 Laplace's equation in spherical polar coordinates

4.4.2 Legendre Polynomials

4.4.3 Solution to radial part

4.5 Multipole expansions for Laplace's equation

4.6 Laplace's equation in cylindrical coordinates

4.7 The heat equation

Proposition. Suppose $\phi : \Omega \times [0, \infty) \rightarrow \mathbb{R}$ satisfies the heat equation $\frac{\partial\phi}{\partial t} = \kappa\nabla^2\phi$, and obeys

- Initial conditions $\phi(\mathbf{x}, 0) = f(x)$ for all $x \in \Omega$
- Boundary condition $\phi(\mathbf{x}, t)|_{\partial\Omega} = g(\mathbf{x}, t)$ for all $t \in [0, \infty)$.

Then $\phi(\mathbf{x}, t)$ is unique.

4.8 The wave equation

Proposition. Suppose $\phi : \Omega \times [0, \infty) \rightarrow \mathbb{R}$ obeys the wave equation $\frac{\partial^2\phi}{\partial t^2} = c^2\nabla^2\phi$ inside $\Omega \times (0, \infty)$, and is fixed at the boundary. Then E is constant.

Proposition. Suppose $\phi : \Omega \times [0, \infty) \rightarrow \mathbb{R}$ obeys the wave equation $\frac{\partial^2\phi}{\partial t^2} = c^2\nabla^2\phi$ inside $\Omega \times (0, \infty)$, and obeys, for some f, g, h ,

- (i) $\phi(x, 0) = f(x)$;
- (ii) $\frac{\partial\phi}{\partial t}(x, 0) = g(x)$; and
- (iii) $\phi|_{\partial\Omega \times [0, \infty)} = h(x)$.

Then ϕ is unique.

5 Distributions

5.1 Distributions

5.2 Green's functions

5.3 Green's functions for initial value problems

6 Fourier transforms

6.1 The Fourier transform

6.2 The Fourier inversion theorem

6.3 Parseval's theorem for Fourier transforms

Theorem (Parseval's theorem (again)). Suppose $f, g : \mathbb{R} \rightarrow \mathbb{C}$ are sufficiently well-behaved that \tilde{f} and \tilde{g} exist and we indeed have $\mathcal{F}^{-1}[\tilde{f}] = f, \mathcal{F}^{-1}[\tilde{g}] = g$. Then

$$(f, g) = \int_{\mathbb{R}} f^*(x)g(x) dx = \frac{1}{2\pi}(\tilde{f}, \tilde{g}).$$

In particular, if $f = g$, then

$$\|f\|^2 = \frac{1}{2\pi}\|\tilde{f}\|^2.$$

6.4 A word of warning

6.5 Fourier transformation of distributions

6.6 Linear systems and response functions

6.7 General form of transfer function

6.8 The discrete Fourier transform

6.9 The fast Fourier transform*

7 More partial differential equations

7.1 Well-posedness

7.2 Method of characteristics

7.3 Characteristics for 2nd order partial differential equations

7.4 Green's functions for PDEs on \mathbb{R}^n

7.5 Poisson's equation

Proposition (Green's first identity).

$$\int_{\partial\Omega} \phi \mathbf{n} \cdot \nabla \psi \, dS = \int_{\Omega} \phi \nabla^2 \psi + (\nabla \phi) \cdot (\nabla \psi) \, dV.$$

Proposition (Green's second identity).

$$\int_{\Omega} \phi \nabla^2 \psi - \psi \nabla^2 \phi \, dV = \int_{\partial\Omega} \phi \mathbf{n} \cdot \nabla \psi - \psi \mathbf{n} \cdot \nabla \phi \, dS.$$

Proposition (Green's third identity).

$$\phi(\mathbf{y}) = \int_{\partial\Omega} \phi(\mathbf{n} \cdot \nabla G_3) - G_3(\mathbf{n} \cdot \nabla \phi) \, dS - \int_{\Omega} G_3(\mathbf{x}, \mathbf{y}) F(\mathbf{x}) \, d^3x.$$