

Part III — Theoretical Physics of Soft Condensed Matter Theorems

Based on lectures by M. E. Cates

Notes taken by Dexter Chua

Lent 2018

These notes are not endorsed by the lecturers, and I have modified them (often significantly) after lectures. They are nowhere near accurate representations of what was actually lectured, and in particular, all errors are almost surely mine.

Soft Condensed Matter refers to liquid crystals, emulsions, molten polymers and other microstructured fluids or semi-solid materials. Alongside many high-tech examples, domestic and biological instances include mayonnaise, toothpaste, engine oil, shaving cream, and the lubricant that stops our joints scraping together. Their behaviour is classical ($\hbar = 0$) but rarely is it deterministic: thermal noise is generally important.

The basic modelling approach therefore involves continuous classical field theories, generally with noise so that the equations of motion are stochastic PDEs. The form of these equations is helpfully constrained by the requirement that the Boltzmann distribution is regained in the steady state (when this indeed holds, i.e. for systems in contact with a heat bath but not subject to forcing). Both the dynamical and steady-state behaviours have a natural expression in terms of path integrals, defined as weighted sums of trajectories (for dynamics) or configurations (for steady state). These concepts will be introduced in a relatively informal way, focusing on how they can be used for actual calculations.

In many cases mean-field treatments are sufficient, simplifying matters considerably. But we will also meet examples such as the phase transition from an isotropic fluid to a ‘smectic liquid crystal’ (a layered state which is periodic, with solid-like order, in one direction but can flow freely in the other two). Here mean-field theory gets the wrong answer for the order of the transition, but the right one is found in a self-consistent treatment that lies one step beyond mean-field (and several steps short of the renormalization group, whose application to classical field theories is discussed in other courses but not this one).

Important models of soft matter include diffusive ϕ^4 field theory (‘Model B’), and the noisy Navier–Stokes equation which describes fluid mechanics at colloidal scales, where the noise term is responsible for Brownian motion of suspended particles in a fluid. Coupling these together creates ‘Model H’, a theory that describes the physics of fluid-fluid mixtures (that is, emulsions). We will explore Model B, and then Model H, in some depth. We will also explore the continuum theory of nematic liquid crystals, which spontaneously break rotational but not translational symmetry, focusing on topological defects and their associated mathematical structure such as homotopy classes.

Finally, the course will cover some recent extensions of the same general approach to systems whose microscopic dynamics does not have time-reversal symmetry, such as self-propelled colloidal swimmers. These systems do not have a Boltzmann distribution in steady state; without that constraint, new field theories arise that are the subject of ongoing research.

Pre-requisites

Knowledge of Statistical Mechanics at an undergraduate level is essential. This course complements the following Michaelmas Term courses although none are prerequisites: Statistical Field Theory; Biological Physics and Complex Fluids; Slow Viscous Flow; Quantum Field Theory.

Contents

0	Introduction	4
0.1	The physics	4
0.2	The mathematics	4
1	Revision of equilibrium statistical physics	5
1.1	Thermodynamics	5
1.2	Coarse Graining	5
2	Mean field theory	6
2.1	Binary fluids	6
2.2	Nematic liquid crystals	6
3	Functional derivatives and integrals	7
3.1	Functional derivatives	7
3.2	Functional integrals	7
4	The variational method	8
4.1	The variational method	8
4.2	Smectic liquid crystals	8
5	Dynamics	9
5.1	A single particle	9
5.2	The Fokker–Planck equation	9
5.3	Field theories	9
6	Model B	10
7	Model H	11
8	Liquid crystals hydrodynamics	12
8.1	Liquid crystal models	12
8.2	Coarsening dynamics for nematics	12
8.3	Topological defects in three dimensions	12
9	Active Soft Matter	13

0 Introduction

0.1 The physics

0.2 The mathematics

1 Revision of equilibrium statistical physics

1.1 Thermodynamics

1.2 Coarse Graining

2 Mean field theory

2.1 Binary fluids

2.2 Nematic liquid crystals

3 Functional derivatives and integrals

3.1 Functional derivatives

3.2 Functional integrals

4 The variational method

4.1 The variational method

4.2 Smectic liquid crystals

5 Dynamics

5.1 A single particle

5.2 The Fokker–Planck equation

5.3 Field theories

6 **Model B**

7 Model H

8 Liquid crystals hydrodynamics

8.1 Liquid crystal models

8.2 Coarsening dynamics for nematics

8.3 Topological defects in three dimensions

9 Active Soft Matter