

Part II — Linear Analysis

Based on lectures by J. W. Luk

Notes taken by Dexter Chua

Michaelmas 2015

These notes are not endorsed by the lecturers, and I have modified them (often significantly) after lectures. They are nowhere near accurate representations of what was actually lectured, and in particular, all errors are almost surely mine.

Part IB Linear Algebra, Analysis II and Metric and Topological Spaces are essential

Normed and Banach spaces. Linear mappings, continuity, boundedness, and norms. Finite-dimensional normed spaces. [4]

The Baire category theorem. The principle of uniform boundedness, the closed graph theorem and the inversion theorem; other applications. [5]

The normality of compact Hausdorff spaces. Urysohn's lemma and Tietze's extension theorem. Spaces of continuous functions. The Stone-Weierstrass theorem and applications. Equicontinuity: the Ascoli-Arzelà theorem. [5]

Inner product spaces and Hilbert spaces; examples and elementary properties. Orthonormal systems, and the orthogonalization process. Bessel's inequality, the Parseval equation, and the Riesz-Fischer theorem. Duality; the self duality of Hilbert space. [5]

Bounded linear operations, invariant subspaces, eigenvectors; the spectrum and resolvent set. Compact operators on Hilbert space; discreteness of spectrum. Spectral theorem for compact Hermitian operators. [5]

Contents

| | | |
|----------|---|-----------|
| 0 | Introduction | 3 |
| 1 | Normed vector spaces | 4 |
| 1.1 | Bounded linear maps | 8 |
| 1.2 | Dual spaces | 9 |
| 1.3 | Adjoint | 10 |
| 1.4 | The double dual | 11 |
| 1.5 | Isomorphism | 12 |
| 1.6 | Finite-dimensional normed vector spaces | 13 |
| 1.7 | Hahn–Banach Theorem | 17 |
| 2 | Baire category theorem | 23 |
| 2.1 | The Baire category theorem | 23 |
| 2.2 | Some applications | 24 |
| 3 | The topology of $C(K)$ | 32 |
| 3.1 | Normality of compact Hausdorff spaces | 32 |
| 3.2 | Tietze-Urysohn extension theorem | 34 |
| 3.3 | Arzelà-Ascoli theorem | 37 |
| 3.4 | Stone–Weierstrass theorem | 41 |
| 4 | Hilbert spaces | 48 |
| 4.1 | Inner product spaces | 48 |
| 4.2 | Riesz representation theorem | 55 |
| 4.3 | Orthonormal systems and basis | 58 |
| 4.4 | The isomorphism with ℓ_2 | 61 |
| 4.5 | Operators | 64 |
| 4.6 | Self-adjoint operators | 76 |

0 Introduction

In IB Linear Algebra, we studied vector spaces in general. Most of the time, we concentrated on finite-dimensional vector spaces, since these are easy to reason about. For example, we know that every finite-dimensional vector space (by definition) has a basis. Using the basis, we can represent vectors and linear maps concretely as column vectors (in \mathbb{F}^n) and matrices.

However, in real life, often we have to work with infinite-dimensional vector spaces instead. For example, we might want to consider the vector space of all continuous (real) functions, or the vector space of infinite sequences. It is difficult to analyse these spaces using the tools from IB Linear Algebra, since many of those assume the vector space is finite-dimensional. Moreover, in these cases, we often are not interested in the vector space structure itself. It's just that the objects we are interested in happen to have a vector space structure. Instead, we want to look at notions like continuity and convergence. We want to do *analysis* on vector spaces. These are not something the vector space structure itself provides.

In this course, we are going to give our vector spaces some additional structure. For the first half of the course, we will grant our vector space a *norm*. This allows us to assign a “length” to each vector. With this, we can easily define convergence and continuity. It turns out this allows us to understand a lot about, say, function spaces and sequence spaces.

In the second half, we will grant a stronger notion, called the inner product. Among many things, this allows us to define orthogonality of the elements of a vector space, which is something we are familiar with from, say, IB Methods.

Most of the time, we will be focusing on infinite-dimensional vector spaces, since finite-dimensional spaces are boring. In fact, we have a section dedicated to proving that finite-dimensional vector spaces are boring. In particular, they are all isomorphic to \mathbb{R}^n , and most of our theorems can be proved trivially for finite-dimensional spaces using what we already know from IB Linear Algebra. So we will not care much about them.

1 Normed vector spaces

In IB Linear Algebra, we have studied vector spaces in quite a lot of detail. However, just knowing something is a vector space usually isn't too helpful. Often, we would want the vector space to have some additional structure. The first structure we will study is a *norm*.

Definition (Normed vector space). A *normed vector space* is a pair $(V, \|\cdot\|)$, where V is a vector space over a field \mathbb{F} and $\|\cdot\|$ is a function $\|\cdot\| : V \mapsto \mathbb{R}$, known as the *norm*, satisfying

- (i) $\|\mathbf{v}\| \geq 0$ for all $\mathbf{v} \in V$, with equality iff $\mathbf{v} = \mathbf{0}$.
- (ii) $\|\lambda\mathbf{v}\| = |\lambda|\|\mathbf{v}\|$ for all $\lambda \in \mathbb{F}, \mathbf{v} \in V$.
- (iii) $\|\mathbf{v} + \mathbf{w}\| \leq \|\mathbf{v}\| + \|\mathbf{w}\|$ for all $\mathbf{v}, \mathbf{w} \in V$.

Intuitively, we think of $\|\mathbf{v}\|$ as the “length” or “magnitude” of the vector.

Example. Let V be a finite dimensional vector space, and $\{e_1, \dots, e_n\}$ a basis. Then, for any $\mathbf{v} = \sum_{i=1}^n v_i \mathbf{e}_i$, we can define a norm as

$$\|\mathbf{v}\| = \sqrt{\sum_{i=1}^n v_i^2}.$$

If we are given a norm of a vector space V , we immediately obtain two more structures on V for free, namely a metric and a topology.

Recall from IB Metric and Topological Spaces that (V, d) is a metric space if the metric $d : V \times V \rightarrow \mathbb{R}$ satisfies

- (i) $d(x, x) = 0$ for all $x \in V$.
- (ii) $d(x, y) = d(y, x)$ for all $x, y \in V$.
- (iii) $d(x, y) \leq d(x, z) + d(z, y)$ for all $x, y, z \in V$.

Also, a topological spaces is a set V together with a topology (a collection of open subsets) such that

- (i) \emptyset and V are open subsets.
- (ii) The union of open subsets is open.
- (iii) The finite intersection of open subsets is open.

As we have seen in IB Metric and Topological Spaces, a norm on a vector space induces a metric by $d(v, w) = \|v - w\|$. This metric in terms defines a topology on V where the open sets are given by “ $U \subseteq V$ is open iff for any $x \in U$, there exists some $\varepsilon > 0$ such that $B(x, \varepsilon) = \{y \in V : d(x, y) < \varepsilon\} \subseteq U$ ”.

This induced topology is not just a random topology on the vector space. They have the nice property that the vector space operators behave well under this topology.

Proposition. Addition $+$: $V \times V \rightarrow V$, and scalar multiplication \cdot : $\mathbb{F} \times V \rightarrow V$ are continuous with respect to the topology induced by the norm (and the usual product topology).

Proof. Let U be open in V . We want to show that $(+)^{-1}(U)$ is open. Let $(\mathbf{v}_1, \mathbf{v}_2) \in (+)^{-1}(U)$, i.e. $\mathbf{v}_1 + \mathbf{v}_2 \in U$. Since $\mathbf{v}_1 + \mathbf{v}_2 \in U$, there exists ε such that $B(\mathbf{v}_1 + \mathbf{v}_2, \varepsilon) \subseteq U$. By the triangle inequality, we know that $B(\mathbf{v}_1, \frac{\varepsilon}{2}) + B(\mathbf{v}_2, \frac{\varepsilon}{2}) \subseteq B(\mathbf{v}_1 + \mathbf{v}_2, \varepsilon) \subseteq U$. Hence we have $(\mathbf{v}_1, \mathbf{v}_2) \in B((\mathbf{v}_1, \mathbf{v}_2), \frac{\varepsilon}{2}) \subseteq (+)^{-1}(U)$. So $(+)^{-1}(U)$ is open.

Scalar multiplication can be done in a very similar way. \square

This motivates the following definition — we can do without the norm, and just require a topology in which addition and scalar multiplication are continuous.

Definition (Topological vector space). A *topological vector space* (V, \mathcal{U}) is a vector space V together with a topology \mathcal{U} such that addition and scalar multiplication are continuous maps, and moreover singleton points $\{\mathbf{v}\}$ are closed sets.

The requirement that points are closed is just a rather technical requirement needed in certain proofs. We should, however, not pay too much attention to this when trying to understand it intuitively.

A natural question to ask is: when is a topological vector space *normable*? i.e. Given a topological vector space, can we find a norm that induces the topology?

To answer this question, we will first need a few definitions.

Definition (Absolute convexity). Let V be a vector space. Then $C \subseteq V$ is *absolutely convex* (or *balanced convex*) if for any $\lambda, \mu \in \mathbb{F}$ such that $|\lambda| + |\mu| \leq 1$, we have $\lambda C + \mu C \subseteq C$. In other words, if $\mathbf{c}_1, \mathbf{c}_2 \in C$, we have $\lambda \mathbf{c}_1 + \mu \mathbf{c}_2 \in C$.

Proposition. If $(V, \|\cdot\|)$ is a normed vector space, then $B(t) = B(\mathbf{0}, t) = \{\mathbf{v} : \|\mathbf{v}\| < t\}$ is absolutely convex.

Proof. By triangle inequality. \square

Definition (Bounded subset). Let V be a topological vector space. Then $B \subseteq V$ is *bounded* if for every open neighbourhood $U \subseteq V$ of $\mathbf{0}$, there is some $s > 0$ such that $B \subseteq tU$ for all $t > s$.

At first sight, this might seem like a rather weird definition. Intuitively, this just means that B is bounded if, whenever we take any open set U , by enlarging it by a scalar multiple, we can make it fully contain B .

Example. $B(t)$ in a normed vector space is bounded.

Proposition. A topological vector space (V, \mathcal{U}) is normable if and only if there exists an absolutely convex, bounded open neighbourhood of $\mathbf{0}$.

Proof. One direction is obvious — if V is normable, then $B(t)$ is an absolutely convex, bounded open neighbourhood of $\mathbf{0}$.

The other direction is not too difficult as well. We define the *Minkowski functional* $\mu : V \rightarrow \mathbb{R}$ by

$$\mu_C(\mathbf{v}) = \inf\{t > 0 : \mathbf{v} \in tC\},$$

where C is our absolutely convex, bounded open neighbourhood.

Note that by definition, for any $t < \mu_C(\mathbf{v})$, $\mathbf{v} \notin tC$. On the other hand, by absolute convexity, for any $t > \mu_C(\mathbf{v})$, we have $\mathbf{v} \in tC$.

We now show that this is a norm on V :

(i) If $\mathbf{v} = \mathbf{0}$, then $\mathbf{v} \in 0C$. So $\mu_C(\mathbf{0}) = 0$. On the other hand, suppose $\mathbf{v} \neq \mathbf{0}$. Since a singleton point is closed, $U = V \setminus \{\mathbf{v}\}$ is an open neighbourhood of $\mathbf{0}$. Hence there is some t such that $C \subseteq tU$. Alternatively, $\frac{1}{t}C \subseteq U$. Hence, $\mathbf{v} \notin \frac{1}{t}C$. So $\mu_C(\mathbf{v}) \geq \frac{1}{t} > 0$. So $\mu_C(\mathbf{v}) = \mathbf{0}$ iff $\mathbf{v} = \mathbf{0}$.

(ii) We have

$$\mu_C(\lambda\mathbf{v}) = \inf\{t > 0 : \lambda\mathbf{v} \in tC\} = \lambda \inf\{t > 0 : \mathbf{v} \in tC\} = \lambda\mu_C(\mathbf{v}).$$

(iii) We want to show that

$$\mu_C(\mathbf{v} + \mathbf{w}) \leq \mu_C(\mathbf{v}) + \mu_C(\mathbf{w}).$$

This is equivalent to showing that

$$\inf\{t > 0 : \mathbf{v} + \mathbf{w} \in tC\} \leq \inf\{t > 0 : \mathbf{v} \in tC\} + \inf\{r > 0 : \mathbf{w} \in rC\}.$$

This is, in turn equivalent to proving that if $\mathbf{v} \in tC$ and $\mathbf{w} \in rC$, then $(\mathbf{v} + \mathbf{w}) \in (t+r)C$.

Let $\mathbf{v}' = \mathbf{v}/t$, $\mathbf{w}' = \mathbf{w}/r$. Then we want to show that if $\mathbf{v}' \in C$ and $\mathbf{w}' \in C$, then $\frac{1}{(t+r)}(t\mathbf{v}' + r\mathbf{w}') \in C$. This is exactly what is required by convexity. So done. \square

In fact, the condition of absolute convexity can be replaced “convex”, where “convex” means for every $t \in [0, 1]$, $tC + (1-t)C \subseteq C$. This is since for every convex bounded C , we can find always find a absolutely convex bounded $\tilde{C} \subseteq C$, which is something not hard to prove.

Among all normed spaces, some are particularly nice, known as Banach spaces.

Definition (Banach spaces). A normed vector space is a *Banach space* if it is complete as a metric space, i.e. every Cauchy sequence converges.

Example.

- (i) A finite dimensional vector space (which is isomorphic to \mathbb{F}^n for some n) is Banach.
- (ii) Let X be a compact Hausdorff space. Then let

$$B(X) = \{f : X \rightarrow \mathbb{R} \text{ such that } f \text{ is bounded}\}.$$

This is obviously a vector space, and we can define the norm be $\|f\| = \sup_{x \in X} f(x)$. It is easy to show that this is a norm. It is less trivial to show that this is a Banach space.

Let $\{f_n\} \subseteq B(X)$ be a Cauchy sequence. Then for any x , $\{f_n(x)\} \subseteq \mathbb{R}$ is also Cauchy. So we can define $f(x) = \lim_{n \rightarrow \infty} f_n(x)$.

To show that $f_n \rightarrow f$, let $\varepsilon > 0$. By definition of f_n being Cauchy, there is some N such that for any $n, m > N$ and any fixed x , we have $|f_n(x) - f_m(x)| < \varepsilon$. Take the limit as $m \rightarrow \infty$. Then $f_m(x) \rightarrow f(x)$. So $|f_n(x) - f(x)| \leq \varepsilon$. Since this is true for all x , for any $n > N$, we must have $\|f_n - f\| \leq \varepsilon$. So $f_n \rightarrow f$.

(iii) Define X as before, and let

$$C(X) = \{f : X \rightarrow \mathbb{R} \text{ such that } f \text{ is continuous}\}.$$

Since any continuous f is bounded, so $C(X) \subseteq B(X)$. We define the norm as before.

Since we know that $C(X) \subseteq B(X)$, to show that $C(X)$ is Banach, it suffices to show that $C(X) \subseteq B(X)$ is closed, i.e. if $f_n \rightarrow f$, $f_n \in C(X)$, then $f \in C(X)$, i.e. the uniform limit of a continuous function is continuous. Proof can be found in IB Analysis II.

(iv) For $1 \leq p < \infty$, define

$$\hat{L}_p([0, 1]) = \{f : [0, 1] \rightarrow \mathbb{R} \text{ such that } f \text{ is continuous}\}.$$

We define the norm $\|\cdot\|_{\hat{L}_p}$ by

$$\|f\|_{\hat{L}_p} = \left(\int_0^1 |f|^p dx \right)^{1/p}.$$

It is easy to show that \hat{L}_p is indeed a vector space, and we now check that this is a norm.

(a) $\|f\|_{\hat{L}_p} \geq 0$ is obvious. Also, suppose that $\|f\|_{\hat{L}_p} = 0$. Then we must have $f = 0$. Otherwise, if $f \neq 0$, say $f(x) = \varepsilon$ for some x . Then there is some δ such that for any $y \in (x - \delta, x + \delta)$, we have $\|f(y)\| \geq \frac{\varepsilon}{2}$. Hence

$$\|f\|_{\hat{L}_p} = \left(\int_0^1 |f|^p dx \right)^{1/p} \geq \left[2\delta \left(\frac{\varepsilon}{2} \right)^p \right]^{1/p} > 0.$$

(b) $\|\lambda f\| = |\lambda| \|f\|$ is obvious

(c) The triangle inequality is the exactly what the Minkowski inequality says, which is in the example sheet.

It turns out that \hat{L}_p is *not* a Banach space. We can brute-force a hard proof here, but we will later develop some tools that allow us to prove this much more easily.

Hence, we define $L_p([0, 1])$ to be the completion of $\hat{L}_p([0, 1])$. In IID Probability and Measure, we will show that $L_p([0, 1])$ is in fact the space

$$L_p([0, 1]) = \left\{ f : [0, 1] \rightarrow \mathbb{R} \text{ such that } \int_0^1 |f|^p dx < \infty \right\} / \sim,$$

where the integral is the Lebesgue integral, and we are quotienting by the relation $f \sim g$ if $f = g$ Lebesgue almost everywhere. You will understand what these terms mean in the IID Probability and Measure course.

(v) ℓ_p spaces: for $p \in [1, \infty)$, define

$$\ell_p(\mathbb{F}) = \left\{ (x_1, x_2, \dots) : x_i \in \mathbb{F}, \sum_{i=1}^{\infty} |x_i|^p < \infty \right\},$$

with the norm

$$\|\mathbf{x}\|_{\ell_p} = \left(\sum_{i=1}^{\infty} |x_i|^p \right)^{1/p}.$$

It should be easy to check that this is a normed vector space. Moreover, this is a Banach space. Proof is in example sheet.

(vi) ℓ_∞ space: we define

$$\ell_\infty = \left\{ (x_1, x_2, \dots) : x_i \in \mathbb{F}, \sup_{i \in \mathbb{N}} |x_i| < \infty \right\}$$

with norm

$$\|\mathbf{x}\|_{\ell_\infty} = \sup_{i \in \mathbb{N}} |x_i|.$$

Again, this is a Banach space.

(vii) Let $B = B(1)$ be the unit open ball in \mathbb{R}^n . Define $C(B)$ to be the set of continuous functions $f : B \rightarrow \mathbb{R}$. Note that unlike in our previous example, these functions need not be bounded. So our previous norm cannot be applied. However, we can still define a topology as follows:

Let $\{K_i\}_{i=1}^{\infty}$ be a sequence of compact subsets of B such that $K_i \subseteq K_{i+1}$ and $\bigcup_{i=1}^{\infty} K_i = B$. We define the basis to include

$$\left\{ f \in C(B) : \sup_{x \in K_i} |f(x)| < \frac{1}{m} \right\}$$

for each $m, i = 1, 2, \dots$, as well as the translations of these sets.

This weird basis is chosen such that $f_n \rightarrow f$ in this topology iff $f_n \rightarrow f$ uniformly in every compact set. It can be showed that this is not normable.

1.1 Bounded linear maps

With vector spaces, we studied linear maps. These are maps that respect the linear structure of a vector space. With *normed* vector spaces, the right kind of maps to study is the *bounded* linear maps.

Definition (Bounded linear map). $T : X \rightarrow Y$ is a *bounded linear map* if there is a constant $C > 0$ such that $\|T\mathbf{x}\|_Y \leq C\|\mathbf{x}\|_X$ for all $\mathbf{x} \in X$. We write $\mathcal{B}(X, Y)$ for the set of bounded linear maps from X to Y .

This is equivalent to saying $T(B_X(1)) \subseteq B_Y(C)$ for some $C > 0$. This also equivalent to saying that $T(B)$ is bounded for every bounded subset B of X . Note that this final characterization is also valid when we just have a topological vector space.

How does boundedness relate to the topological structure of the vector spaces? It turns out that boundedness is the same as continuity, which is another reason why we like bounded linear maps.

Proposition. Let X, Y be normed vector spaces, $T : X \rightarrow Y$ a linear map. Then the following are equivalent:

- (i) T is continuous.
- (ii) T is continuous at $\mathbf{0}$.
- (iii) T is bounded.

Proof. (i) \Rightarrow (ii) is obvious.

(ii) \Rightarrow (iii): Consider $B_Y(1) \subseteq Y$, the unit open ball. Since T is continuous at $\mathbf{0}$, $T^{-1}(B_Y(1)) \subseteq X$ is open. Hence there exists $\varepsilon > 0$ such that $B_X(\varepsilon) \subseteq T^{-1}(B_Y(1))$. So $T(B_X(\varepsilon)) \subseteq B_Y(1)$. So $T(B_X(1)) \subseteq B_Y(\frac{1}{\varepsilon})$. So T is bounded.

(iii) \Rightarrow (i): Let $\varepsilon > 0$. Then $\|T\mathbf{x}_1 - T\mathbf{x}_2\|_Y = \|T(\mathbf{x}_1 - \mathbf{x}_2)\|_Y \leq C\|\mathbf{x}_1 - \mathbf{x}_2\|_X$. This is less than ε if $\|\mathbf{x}_1 - \mathbf{x}_2\|_X < C^{-1}\varepsilon$. So done. \square

Using the obvious operations, $\mathcal{B}(X, Y)$ can be made a vector space. What about a norm?

Definition (Norm on $\mathcal{B}(X, Y)$). Let $T : X \rightarrow Y$ be a bounded linear map. Define $\|T\|_{\mathcal{B}(X, Y)}$ by

$$\|T\|_{\mathcal{B}(X, Y)} = \sup_{\|\mathbf{x}\| \leq 1} \|T\mathbf{x}\|_Y.$$

Alternatively, this is the minimum C such that $\|T\mathbf{x}\|_Y \leq C\|\mathbf{x}\|_X$ for all \mathbf{x} . In particular, we have

$$\|T\mathbf{x}\|_Y \leq \|T\|_{\mathcal{B}(X, Y)}\|\mathbf{x}\|_X.$$

1.2 Dual spaces

We will frequently be interested in one particular case of $\mathcal{B}(X, Y)$.

Definition (Dual space). Let V be a normed vector space. The *dual space* is

$$V^* = \mathcal{B}(V, \mathbb{F}).$$

We call the elements of V^* *functionals*. The *algebraic dual* of V is

$$V' = \mathcal{L}(V, \mathbb{F}),$$

where we do not require boundedness.

One particularly nice property of the dual is that V^* is always a Banach space.

Proposition. Let V be a normed vector space. Then V^* is a Banach space.

Proof. Suppose $\{T_i\} \in V^*$ is a Cauchy sequence. We define T as follows: for any $\mathbf{v} \in V$, $\{T_i(\mathbf{v})\} \subseteq \mathbb{F}$ is Cauchy sequence. Since \mathbb{F} is complete (it is either \mathbb{R} or \mathbb{C}), we can define $T : V \rightarrow \mathbb{F}$ by

$$T(\mathbf{v}) = \lim_{n \rightarrow \infty} T_n(\mathbf{v}).$$

Our objective is to show that $T_i \rightarrow T$. The first step is to show that we indeed have $T \in V^*$, i.e. T is a bounded map.

Let $\|\mathbf{v}\| \leq 1$. Pick $\varepsilon = 1$. Then there is some N such that for all $i > N$, we have

$$|T_i(\mathbf{v}) - T(\mathbf{v})| < 1.$$

Then we have

$$\begin{aligned} |T(\mathbf{v})| &\leq |T_i(\mathbf{v}) - T(\mathbf{v})| + |T_i(\mathbf{v})| \\ &< 1 + \|T_i\|_{V^*} \|\mathbf{v}\|_V \\ &\leq 1 + \|T_i\|_{V^*} \\ &\leq 1 + \sup_i \|T_i\|_{V^*} \end{aligned}$$

Since T_i is Cauchy, $\sup_i \|T_i\|_{V^*}$ is bounded. Since this bound does not depend on \mathbf{v} (and N), we get that T is bounded.

Now we want to show that $\|T_i - T\|_{V^*} \rightarrow 0$ as $n \rightarrow \infty$.

For arbitrary $\varepsilon > 0$, there is some N such that for all $i, j > N$, we have

$$\|T_i - T_j\|_{V^*} < \varepsilon.$$

In particular, for any \mathbf{v} such that $\|\mathbf{v}\| \leq 1$, we have

$$|T_i(\mathbf{v}) - T_j(\mathbf{v})| < \varepsilon.$$

Taking the limit as $j \rightarrow \infty$, we obtain

$$|T_i(\mathbf{v}) - T(\mathbf{v})| \leq \varepsilon.$$

Since this is true for any \mathbf{v} , we have

$$\|T_i - T\|_{V^*} \leq \varepsilon.$$

for all $i > N$. So $T_i \rightarrow T$. □

Exercise: in general, for X, Y normed vector spaces, what condition on X and Y guarantees that $\mathcal{B}(X, Y)$ is a Banach space?

1.3 Adjoint

The idea of the adjoint is given a $T \in \mathcal{B}(X, Y)$, produce a “dual map”, or an *adjoint* $T^* \in \mathcal{B}(Y^*, X^*)$.

There is really only one (non-trivial) natural way of doing this. First we can think about what T^* should do. It takes in something from Y^* and produces something in X^* . By the definition of the dual space, this is equivalent to taking in a function $g : Y \rightarrow \mathbb{F}$ and returning a function $T^*(g) : X \rightarrow \mathbb{F}$.

To produce this $T^*(g)$, the only things we have on our hands to use are $T : X \rightarrow Y$ and $g : Y \rightarrow \mathbb{F}$. Thus the only option we have is to define $T^*(g)$ as the composition $g \circ T$, or $T^*(g)(\mathbf{x}) = g(T(\mathbf{x}))$ (we also have a silly option of producing the zero map regardless of input, but this is silly). Indeed, this is the definition of the adjoint.

Definition (Adjoint). Let X, Y be normed vector spaces. Given $T \in \mathcal{B}(X, Y)$, we define the *adjoint* of T , denoted T^* , as a map $T^* \in \mathcal{B}(Y^*, X^*)$ given by

$$T^*(g)(\mathbf{x}) = g(T(\mathbf{x}))$$

for $\mathbf{x} \in X$, $g \in Y^*$. Alternatively, we can write

$$T^*(g) = g \circ T.$$

It is easy to show that our T^* is indeed linear. We now show it is bounded.

Proposition. T^* is bounded.

Proof. We want to show that $\|T^*\|_{\mathcal{B}(Y^*, X^*)}$ is finite. For simplicity of notation, the supremum is assumed to be taken over non-zero elements of the space. We have

$$\begin{aligned} \|T^*\|_{\mathcal{B}(Y^*, X^*)} &= \sup_{g \in Y^*} \frac{\|T^*(g)\|_{X^*}}{\|g\|_{Y^*}} \\ &= \sup_{g \in Y^*} \sup_{\mathbf{x} \in X} \frac{|T^*(g)(\mathbf{x})| / \|\mathbf{x}\|_X}{\|g\|_{Y^*}} \\ &= \sup_{g \in Y^*} \sup_{\mathbf{x} \in X} \frac{|g(T\mathbf{x})|}{\|g\|_{Y^*} \|\mathbf{x}\|_X} \\ &\leq \sup_{g \in Y^*} \sup_{\mathbf{x} \in X} \frac{\|g\|_{Y^*} \|T\mathbf{x}\|_Y}{\|g\|_{Y^*} \|\mathbf{x}\|_X} \\ &\leq \sup_{\mathbf{x} \in X} \frac{\|T\|_{\mathcal{B}(X, Y)} \|\mathbf{x}\|_X}{\|\mathbf{x}\|_X} \\ &= \|T\|_{\mathcal{B}(X, Y)} \end{aligned}$$

So it is finite. □

1.4 The double dual

Definition (Double dual). Let V be a normed vector space. Define $V^{**} = (V^*)^*$.

We want to define a map $\phi : V \rightarrow V^{**}$. Again, we can reason about what we expect this function to do. It takes in a $\mathbf{v} \in V$, and produces a $\phi(\mathbf{v}) \in V^{**}$. Expanding the definition, this gives a $\phi(\mathbf{v}) : V^* \rightarrow \mathbb{F}$. Hence this $\phi(\mathbf{v})$ takes in a $g \in V^*$, and returns a $\phi(\mathbf{v})(g) \in \mathbb{F}$.

This is easy. Since $g \in V^*$, we know that g is a function $g : V \rightarrow \mathbb{F}$. Given this function g and a $\mathbf{v} \in V$, it is easy to produce a $\phi(\mathbf{v})(g) \in \mathbb{F}$. Just apply g on \mathbf{v} :

$$\phi(\mathbf{v})(g) = g(\mathbf{v}).$$

Proposition. Let $\phi : V \rightarrow V^{**}$ be defined by $\phi(\mathbf{v})(g) = g(\mathbf{v})$. Then ϕ is a bounded linear map and $\|\phi\|_{\mathcal{B}(V, V^{**})} \leq 1$

Proof. Again, we are taking supremum over non-zero elements. We have

$$\begin{aligned} \|\phi\|_{\mathcal{B}(V, V^{**})} &= \sup_{\mathbf{v} \in V} \frac{\|\phi(\mathbf{v})\|_{V^{**}}}{\|\mathbf{v}\|_V} \\ &= \sup_{\mathbf{v} \in V} \sup_{g \in V^*} \frac{|\phi(\mathbf{v})(g)|}{\|\mathbf{v}\|_V \|g\|_{V^*}} \\ &= \sup_{\mathbf{v} \in V} \sup_{g \in V^*} \frac{|g(\mathbf{v})|}{\|\mathbf{v}\|_V \|g\|_{V^*}} \\ &\leq 1. \end{aligned} \quad \square$$

In fact, we will later show that $\|\phi\|_{\mathcal{B}(V, V^{**})} = 1$.

1.5 Isomorphism

So far, we have discussed a lot about bounded linear maps, which are “morphisms” between normed vector spaces. It is thus natural to come up with the notion of isomorphism.

Definition (Isomorphism). Let X, Y be normed vector spaces. Then $T : X \rightarrow Y$ is an *isomorphism* if it is a bounded linear map with a bounded linear inverse (i.e. it is a homeomorphism).

We say X and Y are *isomorphic* if there is an isomorphism $T : X \rightarrow Y$.

We say that $T : X \rightarrow Y$ is an *isometric* isomorphism if T is an isomorphism and $\|T\mathbf{x}\|_Y = \|\mathbf{x}\|_X$ for all $\mathbf{x} \in X$.

X and Y are *isometrically isomorphic* if there is an isometric isomorphism between them.

Example. Consider a finite-dimensional space \mathbb{F}^n with the standard basis $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$. For any $\mathbf{v} = \sum v_i \mathbf{e}_i$, the norm is defined by

$$\|\mathbf{v}\| = \left(\sum v_i^2 \right)^{1/2}.$$

Then any $g \in V^*$ is determined by $g(\mathbf{e}_i)$ for $i = 1, \dots, n$. We want to show that there are no restrictions on what $g(\mathbf{e}_i)$ can be, i.e. whatever values I assign to them, g will still be bounded. We have

$$\begin{aligned} \|g\|_{V^*} &= \sup_{\mathbf{v} \in V} \frac{|g(\mathbf{v})|}{\|\mathbf{v}\|} \\ &\leq \sup_{\mathbf{v} \in V} \frac{\sum |v_i| |g(\mathbf{e}_i)|}{\left(\sum |v_i|^2 \right)^{1/2}} \\ &\leq C \sup_{\mathbf{v} \in V} \frac{\left(\sum |v_i|^2 \right)^{1/2}}{\left(\sum |v_i|^2 \right)^{1/2}} \left(\sup_i |g(\mathbf{e}_i)| \right) \\ &= C \sup_i |g(\mathbf{e}_i)| \end{aligned}$$

for some C , where the second-to-last line is due to the Cauchy-Schwarz inequality.

The supremum is finite since \mathbb{F}^n is finite dimensional.

Since g is uniquely determined by a list of values $(g(\mathbf{e}_1), g(\mathbf{e}_2), \dots, g(\mathbf{e}_n))$, it has dimension n . Therefore, V^* is isomorphic to \mathbb{F}^n . By the same lines of argument, V^{**} is isomorphic to \mathbb{F}^n .

In fact, we can show that $\phi : V \rightarrow V^{**}$ by $\phi(\mathbf{v})(g) = g(\mathbf{v})$ is an isometric isomorphism (this is not true for general normed vector spaces. Just pick V to be incomplete, then V and V^{**} cannot be isomorphic since V^{**} is complete).

Example. Consider ℓ_p for $p \in [1, \infty)$. What is ℓ_p^* ?

Suppose q is the *conjugate exponent* of p , i.e.

$$\frac{1}{q} + \frac{1}{p} = 1.$$

(if $p = 1$, define $q = \infty$) It is easy to see that $\ell_q \subseteq \ell_p^*$ by the following:

Suppose $(x_1, x_2, \dots) \in \ell_p$, and $(y_1, y_2, \dots) \in \ell_q$. Define $y(\mathbf{x}) = \sum_{i=1}^{\infty} x_i y_i$. We will show that y defined this way is a bounded linear map. Linearity is easy

to see, and boundedness comes from the fact that

$$\|y\|_{\ell_p^*} = \sup_{\mathbf{x}} \frac{\|y(\mathbf{x})\|}{\|\mathbf{x}\|_{\ell_p}} = \sup_{\mathbf{x}} \frac{\sum x_i y_i}{\|\mathbf{x}\|_{\ell_p}} \leq \sup_{\mathbf{x}} \frac{\|\mathbf{x}\|_{\ell_p} \|\mathbf{y}\|_{\ell_q}}{\|\mathbf{x}\|_{\ell_p}} = \|\mathbf{y}\|_{\ell_q},$$

by the Hölder's inequality. So every $(y_i) \in \ell_q$ determines a bounded linear map. In fact, we can show ℓ_p^* is isomorphic to ℓ_q .

1.6 Finite-dimensional normed vector spaces

We are now going to look at a special case of normed vector spaces, where the vector space is finite dimensional.

It turns out that finite-dimensional vector spaces are rather boring. In particular, we have

- (i) All norms are equivalent.
- (ii) The closed unit ball is compact.
- (iii) They are Banach spaces.
- (iv) All linear maps whose domain is finite dimensional are bounded.

These are what we are going to show in this section.

First of all, we need to say what we mean when we say all norms are “equivalent”

Definition (Equivalent norms). Let V be a vector space, and $\|\cdot\|_1, \|\cdot\|_2$ be norms on V . We say that these are *equivalent* if there exists a constant $C > 0$ such that for any $\mathbf{v} \in V$, we have

$$C^{-1}\|\mathbf{v}\|_2 \leq \|\mathbf{v}\|_1 \leq C\|\mathbf{v}\|_2.$$

It is an exercise to show that equivalent norms induce the same topology, and hence agree on continuity and convergence. Also, equivalence of norms is an equivalence relation (as the name suggests).

Now let V be an n -dimensional vector space with basis $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$. We can define the ℓ_p^n norm by

$$\|\mathbf{v}\|_{\ell_p^n} = \left(\sum_{i=1}^n |v_i|^p \right)^{1/p},$$

where

$$\mathbf{v} = \sum_{i=1}^n v_i \mathbf{e}_i.$$

Proposition. Let V be an n -dimensional vector space. Then all norms on V are equivalent to the norm $\|\cdot\|_{\ell_1^n}$.

Corollary. All norms on a finite-dimensional vector space are equivalent.

Proof. Let $\|\cdot\|$ be a norm on V .

Let $\mathbf{v} = (v_1, \dots, v_n) = \sum v_i \mathbf{e}_i \in V$. Then we have

$$\begin{aligned} \|\mathbf{v}\| &= \left\| \sum v_i \mathbf{e}_i \right\| \\ &\leq \sum_{i=1}^n |v_i| \|\mathbf{e}_i\| \\ &\leq \left(\sup_i \|\mathbf{e}_i\| \right) \sum_{i=1}^n |v_i| \\ &\leq C \|\mathbf{v}\|_{\ell_1^n}, \end{aligned}$$

where $C = \sup \|\mathbf{e}_i\| < \infty$ since we are taking a finite supremum.

For the other way round, let $S_1 = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\ell_1^n} = 1\}$. We will show the two following results:

- (i) $\|\cdot\| : (S_1, \|\cdot\|_{\ell_1^n}) \rightarrow \mathbb{R}$ is continuous.
- (ii) S_1 is a compact set.

We first see why this gives what we want. We know that for any continuous map from a compact set to \mathbb{R} , the image is bounded and the infimum is achieved. So there is some $\mathbf{v}_* \in S_1$ such that

$$\|\mathbf{v}_*\| = \inf_{\mathbf{v} \in S_1} \|\mathbf{v}\|.$$

Since $\mathbf{v}_* \neq 0$, there is some c' such that $\|\mathbf{v}\| \geq c'$ for all $\mathbf{v} \in S_1$.

Now take an arbitrary non-zero $\mathbf{v} \in V$, since $\frac{\mathbf{v}}{\|\mathbf{v}\|_{\ell_1^n}} \in S_1$, we know that

$$\left\| \frac{\mathbf{v}}{\|\mathbf{v}\|_{\ell_1^n}} \right\| \geq c',$$

which is to say that

$$\|\mathbf{v}\| \geq c' \|\mathbf{v}\|_{\ell_1^n}.$$

Since we have found $c, c' > 0$ such that

$$c' \|\mathbf{v}\|_{\ell_1^n} \leq \|\mathbf{v}\| \leq c \|\mathbf{v}\|_{\ell_1^n},$$

now let $C = \max\{c, \frac{1}{c'}\} > 0$. Then

$$C^{-1} \|\mathbf{v}\|_2 \leq \|\mathbf{v}\|_1 \leq C \|\mathbf{v}\|_2.$$

So the norms are equivalent. Now we can start to prove (i) and (ii).

First, let $\mathbf{v}, \mathbf{w} \in V$. We have

$$\left| \|\mathbf{v}\| - \|\mathbf{w}\| \right| \leq \|\mathbf{v} - \mathbf{w}\| \leq C \|\mathbf{v} - \mathbf{w}\|_{\ell_1^n}.$$

Hence when \mathbf{v} is close to \mathbf{w} under ℓ_1^n , then $\|\mathbf{v}\|$ is close to $\|\mathbf{w}\|$. So it is continuous.

To show (ii), it suffices to show that the unit ball $B = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\ell_1^n} \leq 1\}$ is compact, since S_1 is a closed subset of B . We will do so by showing it is sequentially compact.

Let $\{\mathbf{v}^{(k)}\}_{k=1}^{\infty}$ be a sequence in B . Write

$$\mathbf{v}^{(k)} = \sum_{i=1}^n \lambda_i^{(k)} \mathbf{e}_i.$$

Since $\mathbf{v}^{(k)} \in B$, we have

$$\sum_{i=1}^n |\lambda_i^{(k)}| \leq 1.$$

Consider the sequence $\lambda_1^{(k)}$, which is a sequence in \mathbb{F} .

We know that $|\lambda_1^{(k)}| \leq 1$. So by Bolzano-Weierstrass, there is a convergent subsequence $\lambda_1^{(k_{j_1})}$.

Now look at $\lambda_2^{(k_{j_1})}$. Since this is bounded, there is a convergent subsequence $\lambda_2^{(k_{j_2})}$.

Iterate this for all n to obtain a sequence k_{j_n} such that $\lambda_i^{(k_{j_n})}$ is convergent for all i . So $\mathbf{v}^{(k_{j_n})}$ is a convergent subsequence. \square

Proposition. Let V be a finite-dimensional normed vector space. Then the closed unit ball

$$\bar{B}(1) = \{\mathbf{v} \in V : \|\mathbf{v}\| \leq 1\}$$

is compact.

Proof. This follows from the proof above. \square

Proposition. Let V be a finite-dimensional normed vector space. Then V is a Banach space.

Proof. Let $\{\mathbf{v}_i\} \in V$ be a Cauchy sequence. Since $\{\mathbf{v}_i\}$ is Cauchy, it is bounded, i.e. $\{\mathbf{v}_i\} \subseteq \bar{B}(R)$ for some $R > 0$. By above, $\bar{B}(R)$ is compact. So $\{\mathbf{v}_i\}$ has a convergent subsequence $\mathbf{v}_{i_k} \rightarrow \mathbf{v}$. Since $\{\mathbf{v}_i\}$ is Cauchy, we must have $\mathbf{v}_i \rightarrow \mathbf{v}$. So \mathbf{v}_i converges. \square

Proposition. Let V, W be normed vector spaces, V be finite-dimensional. Also, let $T : V \rightarrow W$ be a linear map. Then T is bounded.

Proof. Recall discussions last time about V^* for finite-dimensional V . We will do a similar proof.

Note that since V is finite-dimensional, $\text{im } T$ finite dimensional. So wlog W is finite-dimensional. Since all norms are equivalent, it suffices to consider the case where the vector spaces have ℓ_1^n and ℓ_1^m norm. This can be represented by a matrix T_{ij} such that

$$T(x_1, \dots, x_n) = \left(\sum T_{1i}x_i, \dots, \sum T_{mi}x_i \right).$$

We can bound this by

$$\|T(x_1, \dots, x_n)\| \leq \sum_{j=1}^m \sum_{i=1}^n |T_{ji}| |x_i| \leq m \left(\sup_{i,j} |T_{ij}| \right) \sum_{i=1}^n |x_i| \leq C \|\mathbf{x}\|_{\ell_1^n}$$

for some $C > 0$, since we are taking the supremum over a finite set. This implies that $\|T\|_{\mathcal{B}(\ell_1^n, \ell_1^m)} \leq C$. \square

There is another way to prove this statement.

Proof. (alternative) Let $T : V \rightarrow W$ be a linear map. We define a norm on V by $\|\mathbf{v}\|' = \|\mathbf{v}\|_V + \|T\mathbf{v}\|_W$. It is easy to show that this is a norm.

Since V is finite dimensional, all norms are equivalent. So there is a constant $C > 0$ such that for all \mathbf{v} , we have

$$\|\mathbf{v}\|' \leq C\|\mathbf{v}\|_V.$$

In particular, we have

$$\|T\mathbf{v}\| \leq C\|\mathbf{v}\|_V.$$

So done. □

Among all these properties, compactness of $\bar{B}(1)$ characterizes finite dimensionality.

Proposition. Let V be a normed vector space. Suppose that the closed unit ball $\bar{B}(1)$ is compact. Then V is finite dimensional.

Proof. Consider the following open cover of $\bar{B}(1)$:

$$\bar{B}(1) \subseteq \bigcup_{\mathbf{y} \in \bar{B}(1)} B\left(\mathbf{y}, \frac{1}{2}\right).$$

Since $\bar{B}(1)$ is compact, this has a finite subcover. So there is some $\mathbf{y}_1, \dots, \mathbf{y}_n$ such that

$$\bar{B}(1) \subseteq \bigcup_{i=1}^n B\left(\mathbf{y}_i, \frac{1}{2}\right).$$

Now let $Y = \text{span}\{\mathbf{y}_1, \dots, \mathbf{y}_n\}$, which is a finite-dimensional subspace of V . We want to show that in fact we have $Y = V$.

Clearly, by definition of Y , the unit ball

$$B(1) \subseteq Y + B\left(\frac{1}{2}\right),$$

i.e. for every $\mathbf{v} \in B(1)$, there is some $\mathbf{y} \in Y, \mathbf{w} \in B(\frac{1}{2})$ such that $\mathbf{v} = \mathbf{y} + \mathbf{w}$. Multiplying everything by $\frac{1}{2}$, we get

$$B\left(\frac{1}{2}\right) \subseteq Y + B\left(\frac{1}{4}\right).$$

Hence we also have

$$B(1) \subseteq Y + B\left(\frac{1}{4}\right).$$

By induction, for every n , we have

$$B(1) \subseteq Y + B\left(\frac{1}{2^n}\right).$$

As a consequence,

$$B(1) \subseteq \bar{Y}.$$

Since Y is finite-dimensional, we know that Y is complete. So Y is a closed subspace of V . So $\bar{Y} = Y$. So in fact

$$B(1) \subseteq Y.$$

Since every element in V can be rescaled to an element of $B(1)$, we know that $V = Y$. Hence V is finite dimensional. \square

This concludes our discussion on finite-dimensional vector spaces. We'll end with an example that shows these are not true for infinite dimensional vector spaces.

Example. Consider ℓ_1 , and $\mathbf{e}_i = (0, 0, \dots, 0, 1, 0, \dots)$, where \mathbf{e}_i is 1 on the i th entry, 0 elsewhere.

Note that if $i \neq j$, then

$$\|\mathbf{e}_i - \mathbf{e}_j\| = 2.$$

Since $\mathbf{e}_i \in \bar{B}(1)$, we see that $\bar{B}(1)$ cannot be covered by finitely many open balls of radius $\frac{1}{2}$, since each open ball can contain at most one of $\{\mathbf{e}_i\}$.

1.7 Hahn–Banach Theorem

Let V be a real normed vector space. What can we say about $V^* = \mathcal{B}(V, \mathbb{R})$? For instance, If V is non-trivial, must V^* be non-trivial?

The main goal of this section is to prove the *Hahn–Banach theorem* (surprise), which allows us to produce a lot of elements in V^* . Moreover, it doesn't just tell us that V^* is non-empty (this is rather dull), but provides a tool to craft (or at least prove existence of) elements of V^* that satisfy some property we want.

Proposition. Let V be a real normed vector space, and $W \subseteq V$ has co-dimension 1. Assume we have the following two items:

– $p : V \rightarrow \mathbb{R}$ (not necessarily linear), which is positive homogeneous, i.e.

$$p(\lambda \mathbf{v}) = \lambda p(\mathbf{v})$$

for all $\mathbf{v} \in V, \lambda > 0$, and subadditive, i.e.

$$p(\mathbf{v}_1 + \mathbf{v}_2) \leq p(\mathbf{v}_1) + p(\mathbf{v}_2)$$

for all $\mathbf{v}_1, \mathbf{v}_2 \in V$. We can think of something like a norm, but more general.

– $f : W \rightarrow \mathbb{R}$ a linear map such that $f(\mathbf{w}) \leq p(\mathbf{w})$ for all $\mathbf{w} \in W$.

Then there exists an extension $\tilde{f} : V \rightarrow \mathbb{R}$ which is linear such that $\tilde{f}|_W = f$ and $\tilde{f}(\mathbf{v}) \leq p(\mathbf{v})$ for all $\mathbf{v} \in V$.

Why do we want this weird theorem? Our objective is to find something in V^* . This theorem tells us that to find a bounded linear map in V , we just need something in W bounded by a norm-like object, and then we can extend it to V .

Proof. Let $\mathbf{v}_0 \in V \setminus W$. Since W has co-dimension 1, every element $\mathbf{v} \in V$ can be written uniquely as $\mathbf{v} = \mathbf{w} + a\mathbf{v}_0$, for some $\mathbf{w} \in W, a \in \mathbb{R}$. Therefore it suffices to define $\tilde{f}(\mathbf{v}_0)$ and then extend linearly to V .

The condition we want to meet is

$$\tilde{f}(\mathbf{w} + a\mathbf{v}_0) \leq p(\mathbf{w} + a\mathbf{v}_0) \quad (*)$$

for all $\mathbf{w} \in W, a \in \mathbb{R}$. If $a = 0$, then this is satisfied since \tilde{f} restricts to f on W .

If $a > 0$ then $(*)$ is equivalent to

$$\tilde{f}(\mathbf{w}) + a\tilde{f}(\mathbf{v}_0) \leq p(\mathbf{w} + a\mathbf{v}_0).$$

We can divide by a to obtain

$$\tilde{f}(a^{-1}\mathbf{w}) + \tilde{f}(\mathbf{v}_0) \leq p(a^{-1}\mathbf{w} + \mathbf{v}_0).$$

We let $\mathbf{w}' = a^{-1}\mathbf{w}$. So we can write this as

$$\tilde{f}(\mathbf{v}_0) \leq p(\mathbf{w}' + \mathbf{v}_0) - f(\mathbf{w}'),$$

for all $\mathbf{w}' \in W$.

If $a < 0$, then $(*)$ is equivalent to

$$\tilde{f}(\mathbf{w}) + a\tilde{f}(\mathbf{v}_0) \leq p(\mathbf{w} + a\mathbf{v}_0).$$

We now divide by a and flip the sign of the equality. So we have

$$\tilde{f}(a^{-1}\mathbf{w}) + \tilde{f}(\mathbf{v}_0) \geq -(-a^{-1})p(\mathbf{w} + a\mathbf{v}_0).$$

In other words, we want

$$\tilde{f}(\mathbf{v}_0) \geq -p(-a^{-1}\mathbf{w} - \mathbf{v}_0) - f(a^{-1}\mathbf{w}).$$

We let $\mathbf{w}' = -a^{-1}\mathbf{w}$. Then we are left with

$$\tilde{f}(\mathbf{v}_0) \geq -p(\mathbf{w}' - \mathbf{v}_0) + f(\mathbf{w}').$$

for all $\mathbf{w}' \in W$.

Hence we are done if we can define a $\tilde{f}(\mathbf{v}_0)$ that satisfies these two conditions. This is possible if and only if

$$-p(\mathbf{w}_1 - \mathbf{v}_0) + f(\mathbf{w}_1) \leq p(\mathbf{w}_2 + \mathbf{v}_0) - f(\mathbf{w}_2)$$

for all $\mathbf{w}_1, \mathbf{w}_2$. This holds since

$$\begin{aligned} f(\mathbf{w}_1) + f(\mathbf{w}_2) &= f(\mathbf{w}_1 + \mathbf{w}_2) \\ &\leq p(\mathbf{w}_1 + \mathbf{w}_2) \\ &= p(\mathbf{w}_1 - \mathbf{v}_0 + \mathbf{w}_2 + \mathbf{v}_0) \\ &\leq p(\mathbf{w}_1 - \mathbf{v}_0) + p(\mathbf{w}_2 + \mathbf{v}_0). \end{aligned}$$

So the result follows. \square

The goal is to “iterate” this to get a similar result without the co-dimension 1 assumption. While we can do this directly for finitely many times, this isn’t helpful (since we already know a lot about finite dimensional normed spaces). To perform an “infinite iteration”, we need the mysterious result known as Zorn’s lemma.

Digression on Zorn's lemma

We first need a few definitions before we can come to Zorn's lemma.

Definition (Partial order). A relation \leq on a set X is a *partial order* if it satisfies

- (i) $x \leq x$ (reflexivity)
- (ii) $x \leq y$ and $y \leq x$ implies $x = y$ (antisymmetry)
- (iii) $x \leq y$ and $y \leq z$ implies $x \leq z$ (transitivity)

Definition (Total order). Let (S, \leq) be a partial order. $T \subseteq S$ is *totally ordered* if for all $x, y \in T$, either $x \leq y$ or $y \leq x$, i.e. every two things are related.

Definition (Upper bound). Let (S, \leq) be a partial order. $S' \subseteq S$ subset. We say $b \in S$ is an *upper bound* of this subset if $x \leq b$ for all $x \in S'$.

Definition (Maximal element). Let (S, \leq) be a partial order. Then $m \in S$ is a *maximal element* if $x \geq m$ implies $x = m$.

The glorious Zorn's lemma tells us that:

Lemma (Zorn's lemma). Let (S, \leq) be a non-empty partially ordered set such that every totally-ordered subset S' has an upper bound in S . Then S has a maximal element.

We will not give a proof of this lemma here, but can explain why it should be true.

We start by picking one element x_0 in S . If it is maximal, then done. Otherwise, there is some $x_1 > x_0$. If this is not maximal, then pick $x_2 > x_1$. We do this to infinity "and beyond" — after picking infinitely many x_i , if we have not yet reached a maximal element, we take an upper bound of this set, and call it x_ω . If this is not maximal, we can continue picking a larger element.

We can do this forever, but if this process never stops, even after infinite time, we would have picked out more elements than there are in S , which is clearly nonsense. Of course, this is hardly a formal proof. The proper proof can be found in the IID Logic and Set Theory course.

Back to vector spaces

The Hahn–Banach theorem is just our previous proposition without the constraint that W has co-dimension 1.

Theorem (Hahn–Banach theorem*). Let V be a real normed vector space, and $W \subseteq V$ a subspace. Assume we have the following two items:

- $p : V \rightarrow \mathbb{R}$ (not necessarily linear), which is positive homogeneous and subadditive;
- $f : W \rightarrow \mathbb{R}$ a linear map such that $f(\mathbf{w}) \leq p(\mathbf{w})$ for all $\mathbf{w} \in W$.

Then there exists an extension $\tilde{f} : V \rightarrow \mathbb{R}$ which is linear such that $\tilde{f}|_W = f$ and $\tilde{f}(\mathbf{v}) \leq p(\mathbf{v})$ for all $\mathbf{v} \in V$.

Proof. Let S be the set of all pairs (\tilde{V}, \tilde{f}) such that

- (i) $W \subseteq \tilde{V} \subseteq V$
- (ii) $\tilde{f} : \tilde{V} \rightarrow \mathbb{R}$ is linear
- (iii) $\tilde{f}|_W = f$
- (iv) $\tilde{f}(\tilde{\mathbf{v}}) \leq p(\tilde{\mathbf{v}})$ for all $\tilde{\mathbf{v}} \in \tilde{V}$

We introduce a partial order \leq on S by $(\tilde{V}_1, \tilde{f}_1) \leq (\tilde{V}_2, \tilde{f}_2)$ if $\tilde{V}_1 \subseteq \tilde{V}_2$ and $\tilde{f}_2|_{\tilde{V}_1} = \tilde{f}_1$. It is easy to see that this is indeed a partial order.

We now check that this satisfies the assumptions of Zorn's lemma. Let $\{(\tilde{V}_\alpha, \tilde{f}_\alpha)\}_{\alpha \in A} \subseteq S$ be a totally ordered set. Define (\tilde{V}, \tilde{f}) by

$$\tilde{V} = \bigcup_{\alpha \in A} \tilde{V}_\alpha, \quad \tilde{f}(\mathbf{x}) = \tilde{f}_\alpha(\mathbf{x}) \text{ for } \mathbf{x} \in \tilde{V}_\alpha.$$

This is well-defined because $\{(\tilde{V}, \tilde{f}_\alpha)\}_{\alpha \in A}$ is totally ordered. So if $\mathbf{x} \in \tilde{V}_{\alpha_1}$ and $\mathbf{x} \in \tilde{V}_{\alpha_2}$, wlog assume $(\tilde{V}_{\alpha_1}, \tilde{f}_{\alpha_1}) \leq (\tilde{V}_{\alpha_2}, \tilde{f}_{\alpha_2})$. So $\tilde{f}_{\alpha_2}|_{\tilde{V}_{\alpha_1}} = \tilde{f}_{\alpha_1}$. So $\tilde{f}_{\alpha_1}(\mathbf{x}) = \tilde{f}_{\alpha_2}(\mathbf{x})$.

It should be clear that $(\tilde{V}, \tilde{f}) \in S$ and (\tilde{V}, \tilde{f}) is indeed an upper bound of $\{(\tilde{V}_\alpha, \tilde{f}_\alpha)\}_{\alpha \in A}$. So the conditions of Zorn's lemma are satisfied.

Hence by Zorn's lemma, there is a maximal element $(\tilde{W}, \tilde{f}) \in S$. Then by definition, \tilde{f} is linear, restricts to f on W , and bounded by p . We now show that $\tilde{W} = V$.

Suppose not. Then there is some $\mathbf{v}_0 \in V \setminus \tilde{W}$. Define $\tilde{V} = \text{span}\{\tilde{W}, \mathbf{v}_0\}$. Now \tilde{W} is a co-dimensional 1 subspace of \tilde{V} . By our previous result, we know that there is some $\tilde{f} : \tilde{V} \rightarrow \mathbb{R}$ linear such that $\tilde{f}|_{\tilde{W}} = \tilde{f}$ and $\tilde{f}(\mathbf{v}) \leq p(\mathbf{v})$ for all $\mathbf{v} \in \tilde{V}$.

Hence we have $(\tilde{W}, \tilde{f}) \in S$ but $(\tilde{W}, \tilde{f}) < (\tilde{V}, \tilde{f})$. This contradicts the maximality of (\tilde{W}, \tilde{f}) . \square

There is a particularly important special case of this, which is also known as Hahn-Banach theorem sometimes.

Corollary (Hahn-Banach theorem 2.0). Let $W \subseteq V$ be real normed vector spaces. Given $f \in W^*$, there exists a $\tilde{f} \in V^*$ such that $\tilde{f}|_W = f$ and $\|\tilde{f}\|_{V^*} = \|f\|_{W^*}$.

Proof. Use the Hahn-Banach theorem with $p(\mathbf{x}) = \|f\|_{W^*} \|\mathbf{x}\|_V$ for all $\mathbf{x} \in V$. Positive homogeneity and subadditivity follow directly from the axioms of the norm. Then by definition $f(\mathbf{w}) \leq p(\mathbf{w})$ for all $\mathbf{w} \in W$. So Hahn-Banach theorem says that there is $\tilde{f} : V \rightarrow \mathbb{R}$ linear such that $\tilde{f}|_W = f$ and $\tilde{f}(\mathbf{v}) \leq p(\mathbf{v}) = \|f\|_{W^*} \|\mathbf{v}\|_V$.

Now notice that

$$\tilde{f}(\mathbf{v}) \leq \|f\|_{W^*} \|\mathbf{v}\|_V, \quad -\tilde{f}(\mathbf{v}) = \tilde{f}(-\mathbf{v}) \leq \|f\|_{W^*} \|\mathbf{v}\|_V$$

implies that $|\tilde{f}(\mathbf{v})| \leq \|f\|_{W^*} \|\mathbf{v}\|_V$ for all $\mathbf{v} \in V$.

On the other hand, we have (again taking supremum over non-zero \mathbf{v})

$$\|\tilde{f}\|_{V^*} = \sup_{\mathbf{v} \in V} \frac{|\tilde{f}(\mathbf{v})|}{\|\mathbf{v}\|_V} \geq \sup_{\mathbf{w} \in W} \frac{|f(\mathbf{w})|}{\|\mathbf{w}\|_W} = \|f\|_{W^*}.$$

So indeed we have $\|\tilde{f}\|_{V^*} = \|f\|_{W^*}$. \square

We'll have some quick corollaries of these theorems.

Proposition. Let V be a real normed vector space. For every $\mathbf{v} \in V \setminus \{0\}$, there is some $f_{\mathbf{v}} \in V^*$ such that $f_{\mathbf{v}}(\mathbf{v}) = \|\mathbf{v}\|_V$ and $\|f_{\mathbf{v}}\|_{V^*} = 1$.

Proof. Apply Hahn-Banach theorem (2.0) with $W = \text{span}\{\mathbf{v}\}$, $f'_{\mathbf{v}}(\mathbf{v}) = \|\mathbf{v}\|_V$. \square

Corollary. Let V be a real normed vector space. Then $\mathbf{v} = \mathbf{0}$ if and only if $f(\mathbf{v}) = 0$ for all $f \in V^*$.

Corollary. Let V be a non-trivial real normed vector space, $\mathbf{v}, \mathbf{w} \in V$ with $\mathbf{v} \neq \mathbf{w}$. Then there is some $f \in V^*$ such that $f(\mathbf{v}) \neq f(\mathbf{w})$.

Corollary. If V is a non-trivial real normed vector space, then V^* is non-trivial.

We now want to restrict the discussion to double duals. We define $\phi : V \rightarrow V^{**}$ as before by $\phi(\mathbf{v})(f) = f(\mathbf{v})$ for $\mathbf{v} \in V, f \in V^*$.

Proposition. The map $\phi : V \rightarrow V^{**}$ is an isometry, i.e. $\|\phi(\mathbf{v})\|_{V^{**}} = \|\mathbf{v}\|_V$.

Proof. We have previously shown that

$$\|\phi\|_{\mathcal{B}(V, V^{**})} \leq 1.$$

It thus suffices to show that the norm is greater than 1, or that

$$\|\phi(\mathbf{v})\|_{V^{**}} \geq \|\mathbf{v}\|_V.$$

We can assume $\mathbf{v} \neq \mathbf{0}$, for which the inequality is trivial. We have

$$\|\phi(\mathbf{v})\|_{V^{**}} = \sup_{f \in V^*} \frac{|\phi(\mathbf{v})(f)|}{\|f\|_{V^*}} \geq \frac{|\phi(\mathbf{v})(f_{\mathbf{v}})|}{\|f_{\mathbf{v}}\|_{V^*}} = |f_{\mathbf{v}}(\mathbf{v})| = \|\mathbf{v}\|_V,$$

where $f_{\mathbf{v}}$ is the function such that $f_{\mathbf{v}}(\mathbf{v}) = \|\mathbf{v}\|_V, \|f_{\mathbf{v}}\|_{V^*} = 1$ as we have previously defined.

So done. \square

In particular, ϕ is injective and one can view ϕ as an isometric embedding of V into V^{**} .

Definition (Reflexive). We say V is *reflexive* if $\phi(V) = V^{**}$.

Note that any reflexive space is Banach, since V^{**} , being the dual of V^* , is reflexive.

You might have heard that for any infinite dimensional vector space V , the dual of V is always strictly larger than V . This does not prevent an infinite dimensional vector space from being reflexive. When we said the dual of V is always strictly larger than V , we are referring to the *algebraic dual*, i.e. the set of all *linear maps* from V to \mathbb{F} . In the definition of reflexive (and everywhere else where we mention “dual” in this course), we mean the *continuous dual*, where we look at the set of all *bounded* linear maps from V to \mathbb{F} . It is indeed possible for the continuous dual to be isomorphic to the original space, even in infinite dimensional spaces, as we will see later.

Example. Finite-dimensional normed vector spaces are reflexive. Also ℓ^p is reflexive for $p \in (1, \infty)$.

Recall that given $T \in \mathcal{B}(V, W)$, we defined $T^* \in \mathcal{B}(W^*, V^*)$ by

$$T^*(f)(\mathbf{v}) = f(T\mathbf{v})$$

for $\mathbf{v} \in V, f \in W^*$.

We have previously shown that

$$\|T^*\|_{\mathcal{B}(W^*, V^*)} \leq \|T\|_{\mathcal{B}(V, W)}.$$

We will now show that in fact equality holds.

Proposition.

$$\|T^*\|_{\mathcal{B}(W^*, V^*)} = \|T\|_{\mathcal{B}(V, W)}.$$

Proof. We have already shown that

$$\|T^*\|_{\mathcal{B}(W^*, V^*)} \leq \|T\|_{\mathcal{B}(V, W)}.$$

For the other inequality, first let $\varepsilon > 0$. Since

$$\|T\|_{\mathcal{B}(V, W)} = \sup_{\mathbf{v} \in V} \frac{\|T\mathbf{v}\|_W}{\|\mathbf{v}\|_V}$$

by definition, there is some $\mathbf{v} \in V$ such that $\|T\mathbf{v}\|_W \geq \|T\|_{\mathcal{B}(V, W)}\|\mathbf{v}\|_V - \varepsilon$. wlog, assume $\|\mathbf{v}\|_V = 1$. So

$$\|T\mathbf{v}\|_W \geq \|T\|_{\mathcal{B}(V, W)} - \varepsilon.$$

Therefore, we get that

$$\begin{aligned} \|T^*\|_{\mathcal{B}(W^*, V^*)} &= \sup_{f \in W^*} \frac{\|T^*(f)\|_{V^*}}{\|f\|_{W^*}} \\ &\geq \|T^*(f_{T\mathbf{v}})\|_{V^*} \\ &\geq |T^*(f_{T\mathbf{v}})(\mathbf{v})| \\ &= |f_{T\mathbf{v}}(T\mathbf{v})| \\ &= \|T\mathbf{v}\|_W \\ &\geq \|T\|_{\mathcal{B}(V, W)} - \varepsilon, \end{aligned}$$

where we used the fact that $\|f_{T\mathbf{v}}\|_{W^*}$ and $\|\mathbf{v}\|_V$ are both 1. Since ε is arbitrary, we are done. \square

2 Baire category theorem

2.1 The Baire category theorem

When we first write the Baire category theorem down, it might seem a bit pointless. However, it turns out to be a really useful result, and we will be able to prove surprisingly many results from it.

In fact, the Baire category theorem itself does not involve normed vector spaces. It works on any metric space. However, most of the applications we have here are about normed vector spaces.

To specify the theorem, we will need some terminology.

Definition (Nowhere dense set). Let X be a topological space. A subset $E \subseteq X$ is *nowhere dense* if \bar{E} has empty interior.

Usually, we will pick E to be closed so that the definition just says that E has an empty interior.

Definition (First/second category, meagre and residual). Let X be a topological space. We say that $Z \subseteq X$ is of *first category* or *meagre* if it is a countable union of nowhere dense sets.

A subset is of *second category* or *non-meagre* if it is not of first category.

A subset is *residual* if $X \setminus Z$ is meagre.

Theorem (Baire category theorem). Let X be a complete metric space. Then X is of second category.

This, by definition, means that it is not a countable union of nowhere dense sets. This is equivalent to saying that if we can write

$$X = \bigcup_{n=1}^{\infty} C_n,$$

where each C_n are closed, then C_n has a non-empty interior for some n .

Alternatively, we can say that if U_n is a countable collection of open dense sets, then $\bigcap_{n=1}^{\infty} U_n \neq \emptyset$ (for if U_n is open dense, then $X \setminus U_n$ is closed with empty interior).

Proof. We will prove that the intersection of a countable collection of open dense sets is non-empty. Let U_n be a countable collection of open dense set.

The key to proving this is completeness, since that is the only information we have. The idea is to construct a sequence, show that it is Cauchy, and prove that the limit is in the intersection.

Construct a sequence $x_n \in X$ and $\varepsilon_n > 0$ as follows: let x_1, ε_1 be defined such that $\overline{B(x_1, \varepsilon_1)} \subseteq U_1$. This exists U_1 is open and dense. By density, there is some $x_1 \in U_1$, and ε_1 exists by openness.

We define the x_n iteratively. Suppose we already have x_n and ε_n . Define $x_{n+1}, \varepsilon_{n+1}$ such that $\overline{B(x_{n+1}, \varepsilon_{n+1})} \subseteq \overline{B(x_n, \varepsilon_n)} \cap U_{n+1}$. Again, this is possible because U_{n+1} is open and dense. Moreover, we choose our ε_{n+1} such that $\varepsilon_{n+1} < \frac{1}{n}$ so that $\varepsilon_n \rightarrow 0$.

Since $\varepsilon_n \rightarrow 0$, we know that x_n is a Cauchy sequence. By completeness of X , we can find an $x \in X$ such that $x_n \rightarrow x$. Since x is the limit of x_n , we know that $x \in \overline{B(x_n, \varepsilon_n)}$ for all n . In particular, $x \in U_n$ for all n . So done. \square

2.2 Some applications

We are going to have a few applications of the Baire category theorem.

Proposition. $\mathbb{R} \setminus \mathbb{Q} \neq \emptyset$, i.e. there is an irrational number.

Of course, we can also prove this directly by, say, showing that $\sqrt{2}$ is irrational, or that noting that \mathbb{R} is uncountable, but \mathbb{Q} is not. However, we can also use the Baire category theorem.

Proof. Recall that we defined \mathbb{R} to be the completion of \mathbb{Q} . So we just have to show that \mathbb{Q} is not complete.

First, note that \mathbb{Q} is countable. Also, for all $q \in \mathbb{Q}$, $\{q\}$ is closed and has empty interior. Hence

$$\mathbb{Q} = \bigcup_{q \in \mathbb{Q}} \{q\}$$

is the countable union of nowhere dense sets. So it is not complete by the Baire category theorem. \square

We will show that there are normed vector spaces which are not Banach spaces.

Proposition. Let $\hat{\ell}_1$ be a normed vector space defined by the vector space

$$V = \{(x_1, x_2, \dots) : x_i \in \mathbb{R}, \exists I \in \mathbb{N} \text{ such that } i > I \Rightarrow x_i = 0\},$$

with componentwise addition and scalar multiplication. This is the space of all sequences that are eventually zero.

We define the norm by

$$\|x\|_{\hat{\ell}_1} = \sum_{i=1}^{\infty} |x_i|.$$

Then $\hat{\ell}_1$ is not a Banach space.

Note that $\hat{\ell}_1$ is not standard notation.

Proof. Let

$$E_n = \{x \in \hat{\ell}_1 : x_i = 0, \forall i \geq n\}.$$

By definition,

$$\hat{\ell}_1 = \bigcup_{n=1}^{\infty} E_n.$$

We now show that E_n is nowhere dense. We first show that E_n is closed. If $x_j \rightarrow x$ in $\hat{\ell}_1$ with $x_j \in E_n$, then since x_j is 0 from the n th component onwards, x is also 0 from the n th component onwards. So we must have $x \in E_n$. So E_n is closed.

We now show that E_n has empty interior. We need to show that for all $x \in E_n$ and $\varepsilon > 0$, there is some $y \in \hat{\ell}_1$ such that $\|y - x\| < \varepsilon$ but $y \notin E_n$. This is also easy. Given $x = (x_1, \dots, x_{n-1}, 0, 0, \dots)$, we consider

$$y = (x_1, \dots, x_{n-1}, \varepsilon/2, 0, 0, \dots).$$

Then $\|y - x\|_{\hat{\ell}_1} < \varepsilon$ but $y \notin E_n$. Hence by the Baire category theorem, $\hat{\ell}_1$ is not complete. \square

Proposition. There exists an $f \in C([0, 1])$ which is nowhere differentiable.

Proof. (sketch) We want to show that the set of all continuous functions which are differentiable at at least one point is contained in a meagre subset of $C([0, 1])$. Then this set cannot be all of $C([0, 1])$ since $C([0, 1])$ is complete.

Let $E_{m,n}$ be the set of all $f \in C([0, 1])$ such that

$$(\exists x)(\forall y) 0 < |y - x| < \frac{1}{m} \Rightarrow |f(y) - f(x)| < n|y - x|.$$

(where the quantifiers range over $[0, 1]$).

We now show that

$$\{f \in C([0, 1]) : f \text{ is differentiable somewhere}\} \subseteq \bigcup_{n,m=1}^{\infty} E_{m,n}.$$

This is easy from definition. Suppose f is differentiable at x_0 . Then by definition,

$$\lim_{y \rightarrow x_0} \frac{f(y) - f(x_0)}{y - x_0} = f'(x_0).$$

Let $n \in \mathbb{N}$ be such that $|f'(x_0)| < n$. Then by definition of the limit, there is some m such that whenever $0 < |y - x| < \frac{1}{m}$, we have $\frac{|f(y) - f(x)|}{|y - x_0|} < n$. So $f \in E_{m,n}$.

Finally, we need to show that each $E_{m,n}$ is closed and has empty interior. This is left as an exercise for the reader. \square

Theorem (Banach-Steinhaus theorem/uniform boundedness principle). Let V be a Banach space and W be a normed vector space. Suppose T_α is a collection of bounded linear maps $T_\alpha : V \rightarrow W$ such that for each fixed $\mathbf{v} \in V$,

$$\sup_{\alpha} \|T_\alpha(\mathbf{v})\|_W < \infty.$$

Then

$$\sup_{\alpha} \|T_\alpha\|_{\mathcal{B}(V,W)} < \infty.$$

This says that if the set of linear maps is pointwise bounded, then they are uniformly bounded.

Proof. Let

$$E_n = \{\mathbf{v} \in V : \sup_{\alpha} \|T_\alpha(\mathbf{v})\|_W \leq n\}.$$

Then by our conditions,

$$V = \bigcup_{n=1}^{\infty} E_n.$$

We can write each E_n as

$$E_n = \bigcap_{\alpha} \{\mathbf{v} \in V : \|T_\alpha(\mathbf{v})\|_W \leq n\}.$$

Since T_α is bounded and hence continuous, so $\{\mathbf{v} \in V : \|T_\alpha(\mathbf{v})\|_W \leq n\}$ is the continuous preimage of a closed set, and is hence closed. So E_n , being the intersection of closed sets, is closed.

By the Baire category theorem, there is some n such that E_n has non-empty interior. In particular, $(\exists n)(\exists \varepsilon > 0)(\exists \mathbf{v}_0 \in V)$ such that for all $\mathbf{v} \in B(\mathbf{v}_0, \varepsilon)$, we have

$$\sup_{\alpha} \|T_{\alpha}(\mathbf{v})\|_W \leq n.$$

Now consider arbitrary $\|\mathbf{v}'\|_V \leq 1$. Then

$$\mathbf{v}_0 + \frac{\varepsilon}{2} \mathbf{v}' \in B(\mathbf{v}_0, \varepsilon).$$

So

$$\sup_{\alpha} \left\| T_{\alpha} \left(\mathbf{v}_0 + \frac{\varepsilon \mathbf{v}'}{2} \right) \right\|_W \leq n.$$


Therefore

$$\sup_{\alpha} \|T_{\alpha} \mathbf{v}'\|_W \leq \frac{2}{\varepsilon} \left(n + \sup_{\alpha} \|T_{\alpha} \mathbf{v}_0\| \right).$$

Note that the right hand side is independent of \mathbf{v}' . So

$$\sup_{\|\mathbf{v}'\| \leq 1} \sup_{\alpha} \|T_{\alpha} \mathbf{v}'\|_W \leq \infty. \quad \square$$

Note that this result is not true for general functions. For example, consider $f : [0, 1] \rightarrow \mathbb{R}$ defined by



Then for all $x \in [0, 1]$, we have

$$\sup_n |f_n(x)| < \infty,$$

but

$$\sup_n \sup_x |f_n(x)| = \infty.$$

However, by a proof very similar to what we had above, we have

Theorem (Osgood). Let $f_n : [0, 1] \rightarrow \mathbb{R}$ be a sequence of continuous functions such that for all $x \in [0, 1]$

$$\sup_n |f_n(x)| < \infty$$

Then there are some a, b with $0 \leq a < b \leq 1$ such that

$$\sup_n \sup_{x \in [a, b]} |f_n(x)| < \infty.$$

Proof. See example sheet. □

Theorem (Open mapping theorem). Let V and W be Banach spaces and $T : V \rightarrow W$ be a bounded surjective linear map. Then T is an open map, i.e. $T(U)$ is an open subset of W whenever U is an open subset of V .

Note that surjectivity is necessary. If $\mathbf{q} \notin T(V)$, then we can scale \mathbf{q} down arbitrarily and still not be in the image of T . So $T(V)$ does not contain an open neighbourhood of $\mathbf{0}$, and hence cannot be open.

Proof. We can break our proof into three parts:

- (i) We first want an easy way to check if a map is an open map. We want to show that T is open if and only if $T(B_V(1)) \supseteq B_W(\varepsilon)$ for some $\varepsilon > 0$. Note that one direction is easy — if T is open, then by definition $T(B_V(1))$ is open, and hence we can find the epsilon required. So we are going to prove the other direction.
- (ii) We show that $\overline{T(B_V(1))} \supseteq B_W(\varepsilon)$ for some $\varepsilon > 0$
- (iii) By rescaling the norm in W , we may wlog the ε obtained above is in fact 1. We then show that if $\overline{T(B_V(1))} \supseteq B_W(1)$, then $T(B_V(1)) \supseteq B_W(\frac{1}{2})$.

We now prove them one by one.

- (i) Suppose $T(B_V(1)) \supseteq B_W(\varepsilon)$ for some $\varepsilon > 0$. Let $U \subseteq V$ be an open set. We want to show that $T(U)$ is open. So let $\mathbf{p} \in U$, $\mathbf{q} = T\mathbf{p}$.

Since U is open, there is some $\delta > 0$ such that $B_V(\mathbf{p}, \delta) \subseteq U$. We can also write the ball as $B_V(\mathbf{p}, \delta) = \mathbf{p} + B_V(\delta)$. Then we have

$$\begin{aligned} T(U) &\supseteq T(\mathbf{p} + B_V(\delta)) \\ &= T\mathbf{p} + T(B_V(\delta)) \\ &= T\mathbf{p} + \delta T(B_V(1)) \\ &\supseteq \mathbf{q} + \delta B_W(\varepsilon) \\ &= \mathbf{q} + B_W(\delta\varepsilon) \\ &= B_W(\mathbf{q}, \delta\varepsilon). \end{aligned}$$

So done.

- (ii) This is the step where we use the Baire category theorem.

Since T is surjective, we can write W as

$$W = \bigcup_{n=1}^{\infty} T(B_V(n)) = \bigcup_{n=1}^{\infty} T(nB_V(1)) = \bigcup_{n=1}^{\infty} \overline{T(nB_V(1))}.$$

We have written W as a countable union of closed sets. Since W is a Banach space, by the Baire category theorem, there is some $n \geq 1$ such that $\overline{T(nB_V(1))}$ has non-empty interior. But since $\overline{T(nB_V(1))} = n\overline{T(B_V(1))}$, and multiplication by n is a homeomorphism, it follows that $\overline{T(B_V(1))}$ has non-empty interior. So there is some $\varepsilon > 0$ and $\mathbf{w}_0 \in W$ such that

$$\overline{T(B_V(1))} \supseteq B_W(\mathbf{w}_0, \varepsilon).$$

We have now found an open ball in the neighbourhood, but we want a ball centered at the origin. We will use linearity in two ways. Firstly, since if $\mathbf{v} \in B_V(1)$, then $-\mathbf{v} \in B_V(1)$. By linearity of T , we know that

$$\overline{T(B_V(1))} \supseteq B_W(-\mathbf{w}_0, \varepsilon).$$

Then by linearity, intuitively, since the image contains the balls $B_W(\mathbf{w}_0, \varepsilon)$ and $B_W(-\mathbf{w}_0, \varepsilon)$, it must contain everything in between. In particular, it must contain $B_W(\varepsilon)$.

To prove this properly, we need some additional work. This is easy if we had $T(B_V(1)) \supseteq B_W(\mathbf{w}_0, \varepsilon)$ instead of the closure of it — for any $\mathbf{w} \in B_W(\varepsilon)$, we let $\mathbf{v}_1, \mathbf{v}_2 \in B_V(1)$ be such that $T(\mathbf{v}_1) = \mathbf{w}_0 + \mathbf{w}$, $T(\mathbf{v}_2) = -\mathbf{w}_0 + \mathbf{w}$. Then $\mathbf{v} = \frac{\mathbf{v}_1 + \mathbf{v}_2}{2}$ satisfies $\|\mathbf{v}\|_V < 1$ and $T(\mathbf{v}) = \mathbf{w}$.

Since we now have the closure instead, we need to mess with sequences. Since $\overline{T(B_V(1))} \supseteq \pm\mathbf{w}_0 + B_W(\varepsilon)$, for any $\mathbf{w} \in B_W(\varepsilon)$, we can find sequences (\mathbf{v}_i) and (\mathbf{u}_i) such that $\|\mathbf{v}_i\|_V, \|\mathbf{u}_i\|_V < 1$ for all i and $T(\mathbf{v}_i) \rightarrow \mathbf{w}_0 + \mathbf{w}$, $T(\mathbf{u}_i) \rightarrow -\mathbf{w}_0 + \mathbf{w}$.

Now by the triangle inequality, we get

$$\left\| \frac{\mathbf{v}_i + \mathbf{u}_i}{2} \right\| < 1,$$

and we also have

$$\frac{\mathbf{v}_i + \mathbf{u}_i}{2} \rightarrow \frac{\mathbf{w}_0 + \mathbf{w}}{2} + \frac{-\mathbf{w}_0 + \mathbf{w}}{2} = \mathbf{w}.$$

So $\mathbf{w} \in \overline{T(B_V(1))}$. So $\overline{T(B_V(1))} \supseteq B_W(\varepsilon)$.

(iii) Let $\mathbf{w} \in B_W(\frac{1}{2})$. For any δ , we know

$$\overline{T(B_V(\delta))} \supseteq B_W(\delta).$$

Thus, picking $\delta = \frac{1}{2}$, we can find some $\mathbf{v}_1 \in V$ such that

$$\|\mathbf{v}_1\|_V < \frac{1}{2}, \quad \|T\mathbf{v}_1 - \mathbf{w}\| < \frac{1}{4}.$$

Suppose we have recursively found \mathbf{v}_n such that

$$\|\mathbf{v}_n\|_V < \frac{1}{2^n}, \quad \|T(\mathbf{v}_1 + \cdots + \mathbf{v}_n) - \mathbf{w}\| < \frac{1}{2^{n+1}}.$$

Then picking $\delta = \frac{1}{2^{n+1}}$, we can find \mathbf{v}_{n+1} satisfying the properties listed above. Then $\sum_{n=1}^{\infty} \mathbf{v}_n$ is Cauchy, hence convergent by completeness. Let \mathbf{v} be the limit. Then

$$\|\mathbf{v}\|_V \leq \sum_{i=1}^{\infty} \|\mathbf{v}_i\|_V < 1.$$

Moreover, by continuity of T , we know $T\mathbf{v} = \mathbf{w}$. So we are done. \square

Note that in this proof, we required both V and W to be Banach spaces. However, we used the completeness in different ways. We used the completeness of V to extract a limit, but we just used the completeness of W to say it is of second category. In particular, it suffices to assume the image of T is of second category, instead of assuming surjectivity. Hence if we know that $T : V \rightarrow W$ is a bounded linear map such that V is Banach and $\text{im} T$ is of second category, then T is open. As a consequence T has to be surjective (its image contains a small open ball which we can scale up arbitrary).

We are now going to look at certain applications of the open mapping theorem

Theorem (Inverse mapping theorem). Let V, W be Banach spaces, and $T : V \rightarrow W$ be a bounded linear map which is both injective and surjective. Then T^{-1} exists and is a bounded linear map.

Proof. We know that T^{-1} as a function of sets exists. It is also easy to show that it is linear since T is linear. By the open mapping theorem, since $T(U)$ is open for all $U \subseteq V$ open. So $(T^{-1})^{-1}(U)$ is open for all $U \subseteq V$. By definition, T^{-1} is continuous. Hence T^{-1} is bounded since boundedness and continuity are equivalent. \square

Theorem (Closed graph theorem). Let V, W be Banach spaces, and $T : V \rightarrow W$ a linear map. If the graph of T is closed, i.e.

$$\Gamma(T) = \{(\mathbf{v}, T(\mathbf{v})) : \mathbf{v} \in V\} \subseteq V \times W$$

is a closed subset of the product space (using the norm $\|(\mathbf{v}, \mathbf{w})\|_{V \times W} = \max\{\|\mathbf{v}\|_V, \|\mathbf{w}\|_W\}$), then T is bounded.

What does this mean? Closedness of the graph means that if $\mathbf{v}_n \rightarrow \mathbf{v}$ in V and $T(\mathbf{v}_n) \rightarrow \mathbf{w}$, then $\mathbf{w} = T(\mathbf{v})$. What we want to show is continuity, which is a stronger statement that if $\mathbf{v}_n \rightarrow \mathbf{v}$, then $T(\mathbf{v}_n)$ converges and converges to $T(\mathbf{v})$.

Proof. Consider $\phi : \Gamma(T) \rightarrow V$ defined by $\phi(\mathbf{v}, T(\mathbf{v})) = \mathbf{v}$. We want to apply the inverse mapping theorem to this. To do so, we need to show a few things. First we need to show that the spaces are Banach spaces. This is easy — $\Gamma(T)$ is a Banach space since it is a closed subset of a complete space, and we are already given that V is Banach.

Now we need to show surjectivity and injectivity. This is surjective since for any $\mathbf{v} \in V$, we have $\phi(\mathbf{v}, T(\mathbf{v})) = \mathbf{v}$. It is also injective since the function T is single-valued.

Finally, we want to show ϕ is bounded. This is since

$$\|\mathbf{v}\|_V \leq \max\{\|\mathbf{v}\|, \|T(\mathbf{v})\|\} = \|(\mathbf{v}, T(\mathbf{v}))\|_{\Gamma(T)}.$$

By the inverse mapping theorem, ϕ^{-1} is bounded, i.e. there is some $C > 0$ such that

$$\max\{\|\mathbf{v}\|_V, \|T(\mathbf{v})\|\} \leq C\|\mathbf{v}\|_V$$

In particular, $\|T(\mathbf{v})\| \leq C\|\mathbf{v}\|_V$. So T is bounded. \square

Example. We define $D(T)$ to be equal to $C^1([0, 1])$ as a vector space, but equipped with the $C([0, 1])$ norm instead. We seek to show that $D(T)$ is not complete.

To do so, consider the differentiation map

$$T : D(T) \rightarrow C([0, 1]),$$

First of all, this is unbounded. Indeed, consider the sequence of functions $f_n(x) = x^n$. Then $\|f_n\|_{C([0,1])} = 1$ for all n . However,

$$\|Tf_n\|_{C([0,1])} = \sup_{x \in [0,1]} nx^{n-1} = n.$$

So T is unbounded.

We claim the graph of T is closed. If so, then since $C([0, 1])$ is complete, the closed graph theorem implies $D(T)$ is not complete.

To check this, suppose we have $f_n \rightarrow f$ in the $C([0, 1])$ norm, and $f'_n \rightarrow g$, again in the $C([0, 1])$ norm. We want to show that $f' = g$.

By the fundamental theorem of calculus, we have

$$f_n(t) = f_n(0) + \int_0^t f'_n(x) \, dx.$$

Hence by uniform convergence of $f'_n \rightarrow g$ and $f_n \rightarrow f$, we have

$$f(t) = f(0) + \int_0^t g(x) \, dx.$$

So by the fundamental theorem of calculus, we know that $f'(t) = g(t)$. So the graph of T is closed.

Since we know that $C([0, 1])$ is complete, this shows that $D(T)$ is not complete.

Example. We can also use the Baire category theorem to understand Fourier series. Let $f : S^1 \rightarrow \mathbb{R}$ be continuous, i.e. $f : [-\pi, \pi] \rightarrow \mathbb{R}$ which is continuous with periodic boundary condition $f(-\pi) = f(\pi)$. We define the Fourier coefficients $\hat{f} : \mathbb{Z} \rightarrow \mathbb{C}$ by

$$\hat{f}(k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ikx} f(x) \, dx.$$

We define the Fourier series by

$$\sum_{k \in \mathbb{Z}} e^{ikz} \hat{f}(k).$$

In particular, we define the partial sums as

$$S_n(f)(x) = \sum_{k=-n}^n e^{ikx} \hat{f}(k).$$

The question we want to ask is: does the Fourier series converge? We are not even asking if it converges back to f . Just if it converges at all. More concretely, we want to know if $S_n(f)(x)$ has a limit as $n \rightarrow \infty$ for f continuous.

Unfortunately, no. We can show that there exists a continuous function f such that $S_n(f)(x)$ diverges. To show this, we can consider $\phi_n : C(S^1) \rightarrow \mathbb{R}$ defined by $\phi_n(f) = S_n(f)(0)$. Assume that

$$\sup_n |\phi_n(f)|$$

is finite for all continuous f . By Banach-Steinhaus theorem, we have

$$\sup_n \|\phi_n\|_{\mathcal{B}(C(S^1), \mathbb{R})} < \infty,$$

On the other hand, we can show that

$$\phi_n(f) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \frac{\sin(n + \frac{1}{2})x}{\sin \frac{x}{2}} dx.$$

It thus suffices to find a sequence f_n such that $\|f_n\|_{C(S^1)} \leq 1$ but

$$\left| \int_{-\pi}^{\pi} f(x) \frac{\sin(n + \frac{1}{2})x}{\sin \frac{x}{2}} dx \right| \rightarrow \infty,$$

which is a contradiction. Details are left for the example sheet.

What's the role of the Banach-Steinhaus theorem here? If we wanted to prove the result directly, then we need to find a *single* function $f \in C(S^1)$ such that $\phi_n(f)$ is unbounded. However, now we just have to find a sequence $f_n \in C(S^1)$ such that $\phi_n(f_n) \rightarrow \infty$. This is much easier.

There is another thing we can ask. Note that if f is continuous, then $|\hat{f}(k)| \rightarrow 0$ as $k \rightarrow \pm\infty$. In fact, this is even true if $f \in L^1(S^1)$, i.e. f is Lebesgue integrable and

$$\int_{-\pi}^{\pi} |f(x)| dx < \infty.$$

A classical question is: do all sequences $\{a_n\} \in \mathbb{C}$ with $|a_n| \rightarrow 0$ as $n \rightarrow \pm\infty$ arise as the Fourier series of some $f \in L^1$? The answer is no. We let \tilde{c}_0 be the set of all such sequences. By the inverse mapping theorem, if the map $\phi : L^1(S^1) \rightarrow \tilde{c}_0$ is surjective, then the inverse is bounded. But this is not true, since we can find a sequence f_n such that $\|f_n\|_{L^1(S^1)} \rightarrow \infty$ but $\sup_k |\hat{f}_k(\ell)| \leq 1$ for all ℓ . Details are again left for the reader.

3 The topology of $C(K)$

Before we start the chapter, it helps to understand the title. In particular, what is $C(K)$? In the chapter, K will denote a compact Hausdorff topological space. We will first define what it means for a space to be Hausdorff.

Definition (Hausdorff space). A topological space X is *Hausdorff* if for all distinct $p, q \in X$, there are $U_p, U_q \subseteq X$ that are open subsets of X such that $p \in U_p$, $q \in U_q$ and $U_p \cap U_q = \emptyset$.

Example. Every metric space is Hausdorff.

What we want to look at here is compact Hausdorff spaces.

Example. $[0, 1]$ is a compact Hausdorff space.

Notation. $C(K)$ is the set of continuous functions $f : K \rightarrow \mathbb{R}$ with the norm

$$\|f\|_{C(K)} = \sup_{x \in K} |f(x)|.$$

There are three themes we will discuss

- (i) There are many functions in $C(K)$. For example, we will show that given a finite set of points $\{p_i\}_{i=1}^n \subseteq K$ and $\{y_i\}_{i=1}^n \subseteq \mathbb{R}$, there is some $f \in C(K)$ such that $f(p_i) = y_i$. We will prove this later. Note that this is trivial for $C([0, 1])$, since we can use piecewise linear functions. However, this is not easy to prove if K is a general compact Hausdorff space. In fact, we can prove a much stronger statement, known as the Tietze-Urysohn theorem.
- (ii) Elements of $C(K)$ can be approximated by nice functions. This should be thought of as a generalization of the Weierstrass approximation theorem, which states that polynomials are dense in $C([0, 1])$, i.e. every continuous function can be approximated uniformly to arbitrary accuracy by polynomials.
- (iii) Compact subsets of $C(K)$. One question we would like to understand is that given a sequence of functions $\{f_k\}_{k=1}^\infty \subseteq C(K)$, when can we extract a convergent subsequence?

3.1 Normality of compact Hausdorff spaces

At this point, it is helpful to introduce a new class of topological spaces.

Definition (Normal space). A topological space X is *normal* if for every disjoint pair of closed subsets C_1, C_2 of X , there exists $U_1, U_2 \subseteq X$ disjoint open such that $C_1 \subseteq U_1, C_2 \subseteq U_2$.

This is similar to being Hausdorff, except that instead of requiring the ability to separate points, we want the ability to separate closed subsets.

In general, one makes the following definition:

Definition (T_i space). A topological space X has the T_1 *property* if for all $x, y \in X$, where $x \neq y$, there exists $U \subseteq X$ open such that $x \in U$ and $y \notin U$.

A topological space X has the T_2 *property* if X is Hausdorff.

A topological space X has the T_3 property if for any $x \in X$, $C \subseteq X$ closed with $x \notin C$, then there are U_x, U_C disjoint open such that $x \in U_x, C \subseteq U_C$. These spaces are called *regular*.

A topological space X has the T_4 property if X is normal.

Note here that T_4 and T_1 together imply T_2 . It suffices to notice that T_1 implies that x is a closed set for all x — for all y , let U_y be such that $y \in U_y$ and $x \notin U_y$. Then we can write

$$X \setminus \{x\} = \bigcup_{y \neq x} U_y,$$

which is open since it is a union of open sets.

More importantly, we have the following theorem:

Theorem. Let X be a Hausdorff space. If $C_1, C_2 \subseteq X$ are *compact* disjoint subsets, then there are some $U_1, U_2 \subseteq X$ disjoint open such that $C_1 \subseteq U_1, C_2 \subseteq U_2$.

In particular, if X is a compact Hausdorff space, then X is normal (since closed subsets of compact spaces are compact).

Proof. Since C_1 and C_2 are disjoint, by the Hausdorff property, for every $p \in C_1$ and $q \in C_2$, there is some $U_{p,q}, V_{p,q} \subseteq X$ disjoint open with $p \in U_{p,q}, q \in V_{p,q}$.

Now fix a p . Then $\bigcup_{q \in C_2} V_{p,q} \supseteq C_2$ is an open cover. Since C_2 is compact, there is a finite subcover, say

$$C_2 \subseteq \bigcup_{i=1}^n V_{p,q_i} \text{ for some } \{q_1, \dots, q_n\} \subseteq C_2.$$

Note that n and q_i depends on which p we picked at the beginning.

Define

$$U_p = \bigcap_{i=1}^n U_{p,q_i}, \quad V_p = \bigcup_{i=1}^n V_{p,q_i}.$$

Since these are finite intersections and unions, U_p and V_p are open. Also, U_p and V_p are disjoint. We also know that $C_2 \subseteq V_p$.

Now note that $\bigcup_{p \in C_1} U_p \supseteq C_1$ is an open cover. By compactness of C_1 , there is a finite subcover, say

$$C_1 \subseteq \bigcup_{j=1}^m U_{p_j} \text{ for some } \{p_1, \dots, p_m\} \subseteq C_1.$$

Now define

$$U = \bigcup_{j=1}^m U_{p_j}, \quad V = \bigcap_{j=1}^m V_{p_j}.$$

Then U and V are disjoint open with $C_1 \subseteq U, C_2 \subseteq V$. So done. \square

Why do we care about this? It turns out it is easier to discuss continuous functions on normal spaces rather than Hausdorff spaces. Hence, if we are given that a space is compact Hausdorff (e.g. $[0, 1]$), then we know it is normal.


3.2 Tietze-Urysohn extension theorem

The objective of this chapter is to show that if we have a continuous function defined on a closed subset of a normal space, then we can extend this to the whole of the space.

We start a special case of this theorem.

Lemma (Urysohn's lemma). Let X be normal and C_0, C_1 be disjoint closed subsets of X . Then there is a $f \in C(X)$ such that $f|_{C_0} = 0$ and $f|_{C_1} = 1$, and $0 \leq f(x) \leq 1$ for all x .

Before we prove this, let's look at a "stupid" example. Let $X = [-1, 2]$. This is compact Hausdorff. We let $C_0 = [-1, 0]$ and $C_1 = [1, 2]$. To construct the function f , we do the obvious thing:



We can define this function f (in $[0, 1]$) by

$$f(x) = \inf \left\{ \frac{a}{2^n} : a, n \in \mathbb{N}, 0 \leq a \leq 2^n, x \leq \frac{a}{2^n} \right\}.$$

This is obviously a rather silly way to write our function out. However, this is what we will end up doing in the proof below. So keep this in mind for now.

Proof. In this proof, all subsets labeled C are closed, and all subsets labeled U are open.

First note that normality is equivalent to the following: suppose $C \subseteq U \subseteq X$, where U is open and C is closed. Then there is some \tilde{C} closed, \tilde{U} open such that $C \subseteq \tilde{U} \subseteq \tilde{C} \subseteq U$.

We start by defining $U_1 = X \setminus C_1$. Since C_0 and C_1 are disjoint, we know that $C_0 \subseteq U_1$. By normality, there exists $C_{\frac{1}{2}}$ and $U_{\frac{1}{2}}$ such that

$$C_0 \subseteq U_{\frac{1}{2}} \subseteq C_{\frac{1}{2}} \subseteq U_1.$$

Then we can find $C_{\frac{1}{4}}, C_{\frac{3}{4}}, U_{\frac{1}{4}}, U_{\frac{3}{4}}$ such that

$$C_0 \subseteq U_{\frac{1}{4}} \subseteq C_{\frac{1}{4}} \subseteq U_{\frac{1}{2}} \subseteq C_{\frac{1}{2}} \subseteq U_{\frac{3}{4}} \subseteq C_{\frac{3}{4}} \subseteq U_1.$$

Iterating this, we get that for all dyadic rationals $q = \frac{a}{2^n}$, $a, n \in \mathbb{N}$, $0 < a < 2^n$, there are some U_q open, C_q closed such that $U_q \subseteq C_q$, with $C_q \subseteq U_{q'}$ if $q < q'$.

We now define f by

$$f(x) = \inf \{ q \in (0, 1] \text{ dyadic rational} : x \in U_q \},$$

with the understanding that $\inf \emptyset = 1$. We now check the properties desired.

- By definition, we have $0 \leq f(x) \leq 1$.
- If $x \in C_0$, then $x \in U_q$ for all q . So $f(x) = 0$.
- If $x \in C_1$, then $x \notin U_q$ for all q . So $f(x) = 1$.

- To show f is continuous, it suffices to check that $\{x : f(x) > \alpha\}$ and $\{x : f(x) < \alpha\}$ are open for all $\alpha \in \mathbb{R}$, as this shows that the pre-images of all open intervals in \mathbb{R} are open. We know that

$$\begin{aligned} f(x) < \alpha &\Leftrightarrow \inf\{q \in (0, 1) \text{ dyadic rational} : x \in U_q\} < \alpha \\ &\Leftrightarrow (\exists q) q < \alpha \text{ and } x \in U_q \\ &\Leftrightarrow x \in \bigcup_{q < \alpha} U_q. \end{aligned}$$

Hence we have

$$\{x : f(x) < \alpha\} = \bigcup_{q < \alpha} U_q.$$

which is open, since each U_q is open for all q . Similarly we know that

$$\begin{aligned} f(x) > \alpha &\Leftrightarrow \inf\{q : x \in U_q\} > \alpha \\ &\Leftrightarrow (\exists q > \alpha) x \notin C_q \\ &\Leftrightarrow x \in \bigcup_{q > \alpha} X \setminus C_q. \end{aligned}$$

Since this is a union of complement of closed sets, this is open. \square

With this, we can already say that there are many continuous functions. We can just pick some values on some C_0, C_1 , and then get a continuous function out of it. However, we can make a stronger statement.

Theorem (Tietze-Urysohn extension theorem). Let X be a normal topological space, and $C \subseteq X$ be a closed subset. Suppose $f : C \rightarrow \mathbb{R}$ is a continuous function. Then there exists an extension $\tilde{f} : X \rightarrow \mathbb{R}$ which is continuous and satisfies $\tilde{f}|_C = f$ and $\|\tilde{f}\|_{C(X)} = \|f\|_{C(C)}$.

This is in some sense similar to the Hahn-Banach theorem, which states that we can extend linear maps to larger spaces.

Note that this implies the Urysohn's lemma, since if C_0 and C_1 are closed, then $C_0 \cup C_1$ is closed. However, we cannot be lazy and not prove Urysohn's lemma, because the proof of this theorem relies on the Urysohn's lemma.

Proof. The idea is to repeatedly use Urysohn's lemma to get better and better approximations. We can assume wlog that $0 \leq f(x) \leq 1$ for all $x \in C$. Otherwise, we just translate and rescale our function. Moreover, we can assume that the $\sup_{x \in C} f(x) = 1$. It suffices to find $\tilde{f} : X \rightarrow \mathbb{R}$ with $\tilde{f}|_C = f$ with $0 \leq \tilde{f}(x) \leq 1$ for all $x \in X$.


We define the sequences of continuous functions $f_i : C \rightarrow \mathbb{R}$ and $g_i : X \rightarrow \mathbb{R}$ for $i \in \mathbb{N}$. We want to think of the sum $\sum_{i=0}^n g_i$ to be the approximations, and f_{n+1} the error on C .

Let $f_0 = f$. This is the error we have when we approximate with the zero function.

We first define g_0 on a subset of X by

$$g_0(x) = \begin{cases} 0 & x \in f_0^{-1}([0, \frac{1}{3}]) \\ \frac{1}{3} & x \in f_0^{-1}([\frac{2}{3}, 1]) \end{cases}.$$

We can then extend this to the whole of X with $0 \leq g_0(x) \leq \frac{1}{3}$ for all x by Urysohn's lemma.



We define

$$f_1 = f_0 - g_0|_C.$$

By construction, we know that $0 \leq f_1 \leq \frac{2}{3}$. This is our first approximation. Note that we have now lowered our maximum error from 1 to $\frac{2}{3}$. We now repeat this.

Given $f_i : C \rightarrow \mathbb{R}$ with $0 \leq f_i \leq \left(\frac{2}{3}\right)^i$, we define g_i by requiring

$$g_i(x) = \begin{cases} 0 & x \in f_i^{-1} \left(\left[0, \frac{1}{3} \left(\frac{2}{3}\right)^i \right] \right) \\ \frac{1}{3} \left(\frac{2}{3}\right)^i & x \in f_i^{-1} \left(\left[\left(\frac{2}{3}\right)^{i+1}, \left(\frac{2}{3}\right)^i \right] \right) \end{cases},$$

and then extending to the whole of X with $0 \leq g_i \leq \frac{1}{3} \left(\frac{2}{3}\right)^i$ and g_i continuous. Again, this exists by Urysohn's lemma. We then define $f_{i+1} = f_i - g_i|_C$.

We then have

$$\sum_{i=0}^n g_i = (f_0 - f_1) + (f_1 - f_2) + \cdots + (f_n - f_{n+1}) = f - f_{n+1}.$$

We also know that

$$0 \leq f_{i+1} \leq \left(\frac{2}{3}\right)^{i+1}.$$

We conclude by letting

$$\tilde{f} = \sum_{i=0}^{\infty} g_i.$$

This exists because we have the bounds

$$0 \leq g_i \leq \frac{1}{3} \left(\frac{2}{3}\right)^i,$$

and hence $\sum_{i=0}^n g_i$ is Cauchy. So the limit exists and is continuous by the completeness of $C(X)$.

Now we check that

$$\sum_{i=0}^n g_i|_C - f = -f_{n+1}.$$

Since we know that $\|f_{n+1}\|_{C(C)} \rightarrow 0$. Therefore, we know that

$$\sum_{i=0}^{\infty} g_i \Big|_C = \tilde{f}|_C = f.$$

Finally, we check the bounds. We need to show that $0 \leq \tilde{f}(x) \leq 1$. This is true since $g_i \geq 0$ for all i , and also

$$|\tilde{f}(x)| \leq \sum_{i=0}^{\infty} g_i(x) \leq \sum_{i=0}^n \frac{1}{3} \left(\frac{2}{3}\right)^i = 1.$$

So done. □

We can already show what was stated last time — if K is compact Hausdorff, $\{p_1, \dots, p_n\} \subseteq K$ a finite set of points, and $\{y_1, \dots, y_n\} \subseteq \mathbb{R}$, then there exists $f : K \rightarrow \mathbb{R}$ continuous such that $f(p_i) = y_i$. This is since compact Hausdorff spaces are normal, and singleton points are closed sets in Hausdorff spaces. In fact, we can prove this directly with the Urysohn's lemma, by, say, requesting functions f_i such that $f_i(p_i) = y_i$, $f_i(p_j) = 0$ for $i \neq j$. Then we just sum all f_i .

Note that normality is necessary for Urysohn's lemma. Since Urysohn's lemma is a special case of the Tietze-Urysohn extension theorem, normality is also necessary for the Tietze-Urysohn extension theorem. In fact, the lemma is *equivalent* to normality. Let C_0, C_1 be disjoint closed sets of X . If there is some $f : X \rightarrow \mathbb{R}$ such that $f|_{C_0} = 0$, $f|_{C_1} = 1$, then $U_0 = f^{-1}\left(\left(-\infty, \frac{1}{3}\right)\right)$ and $U_1 = f^{-1}\left(\left(\frac{2}{3}, \infty\right)\right)$ are open disjoint sets such that $C_0 \subseteq U_0, C_1 \subseteq U_1$.

Closedness of C_0 and C_1 is also necessary in Urysohn's lemma. For example, we cannot extend $f : [0, \frac{1}{2}) \cup (\frac{1}{2}, 1]$ to $[0, 1]$ continuously, where f is defined as

$$f(x) = \begin{cases} 0 & x < \frac{1}{2} \\ 1 & x > \frac{1}{2} \end{cases}.$$

3.3 Arzelà-Ascoli theorem


Let K be compact Hausdorff, and $\{f_n\}_{n=1}^{\infty}$ be a sequence of continuous functions $f_n : K \rightarrow \mathbb{R}$ (or \mathbb{C}). When does (f_n) have a convergent subsequence in $C(K)$? In other words, when is there a subsequence which converges uniformly?

This will be answered by the Arzelà-Ascoli theorem. Before we get to that, we look at some examples.

Example. Let $K = [0, 1]$, and $f_n(x) = n$. This does not have a convergent subsequence in $C(K)$ since it does not even have a subsequence that converges pointwise. This is since f_n is unbounded.

We see that unboundedness is one “enemy” that prevents us from having a convergent subsequence.

Example. We again let $K[0, 1]$, and let f be defined as follows:



We know that f_n does not have a convergent subsequence in $C(K)$, since any subsequence must converge pointwise to

$$f(x) = \begin{cases} 0 & x \neq 0 \\ 1 & x = 0 \end{cases},$$

which is not continuous.

What is happening here? For every n , fixed x and every ε , by continuity of f_n , there is some δ such that $|x - y| < \delta$ implies $|f_n(x) - f_n(y)| < \varepsilon$, but this choice of δ depends on n , and there is no universal choice that works for us. This is another problem that leads to the lack of a limit.

The Arzelà-Ascoli theorem tells us these are the only two “enemies” that prevent us from extracting a convergent subsequence.

To state this theorem, we first need a definition.

Definition (Equicontinuous). Let K be a topological space, and $F \subseteq C(K)$. We say F is *equicontinuous at* $x \in K$ if for every ε , there is some U which is an open neighbourhood of x such that

$$(\forall f \in F)(\forall y \in U) |f(y) - f(x)| < \varepsilon.$$

We say F is *equicontinuous* if it is equicontinuous at x for all $x \in K$.

Theorem (Arzelà-Ascoli theorem). Let K be a compact topological space. Then $F \subseteq C(K)$ is pre-compact, i.e. \bar{F} is compact, if and only if F is bounded and equicontinuous.

This indeed applies to the problem of extracting a uniformly convergent subsequence, since $C(K)$ is a metric space, and compactness is equivalent to sequential compactness. Indeed, let (f_n) be a bounded and equicontinuous sequence in $C(K)$. Then $F = \{f_n : n \in \mathbb{N}\} \subseteq C(K)$ is bounded and equicontinuous. So it is pre-compact, and hence (f_n) , being a sequence in \bar{F} , has a convergent subsequence.

To prove this, it helps to introduce some more terminology and a few lemmas first.

Definition (ε -net). Let X be a metric space, and let $E \subseteq X$. For $\varepsilon > 0$, we say that $N \subseteq X$ is an ε -net for E if and only if $\bigcup_{x \in N} B(x, \varepsilon) \supseteq E$.

Definition (Totally bounded subset). Let X be a metric space, and $E \subseteq X$. We say that E is *totally bounded* for every ε , there is a finite ε -net N_ε for E .

An important result about totally bounded subsets is the following:

Proposition. Let X be a complete metric space. Then $E \subseteq X$ is totally bounded if and only if for every sequence $\{y_i\}_{i=1}^{\infty} \subseteq E$, there is a subsequence which is Cauchy.

By completeness, we can rewrite this as

Corollary. Let X be a complete metric space. Then $E \subseteq X$ is totally bounded if and only if \bar{E} is compact.

We'll prove these later. For now, we assume this corollary and we will prove Arzelà-Ascoli theorem.

Theorem (Arzelà-Ascoli theorem). Let K be a compact topological space. Then $F \subseteq C(K)$ is pre-compact, i.e. \bar{F} is compact, if and only if F is bounded and equicontinuous.

Proof. By the previous corollary, it suffices to prove that F is totally bounded if and only if F is bounded and equicontinuous. We first do the boring direction.

(\Rightarrow) Suppose F is totally bounded. First notice that F is obviously bounded, since F can be written as the finite union of ε -balls, which must be bounded.

Now we show F is equicontinuous. Let $\varepsilon > 0$. Since F is totally bounded, there exists a finite ε -net for F , i.e. there is some $\{f_1, \dots, f_n\} \subseteq F$ such that for every $f \in F$, there exists an $i \in \{1, \dots, n\}$ such that $\|f - f_i\|_{C(K)} < \varepsilon$.

Consider a point $x \in K$. Since $\{f_1, \dots, f_n\}$ are continuous, for each i , there exists a neighbourhood U_i of x such that $|f_i(y) - f_i(x)| < \varepsilon$ for all $y \in U_i$.

Let

$$U = \bigcap_{i=1}^n U_i.$$

Since this is a finite intersection, U is open. Then for any $f \in F$, $y \in U$, we can find some i such that $\|f - f_i\|_{C(K)} < \varepsilon$. So

$$|f(y) - f(x)| \leq |f(y) - f_i(y)| + |f_i(y) - f_i(x)| + |f_i(x) - f(x)| < 3\varepsilon.$$

So F is equicontinuous at x . Since x was arbitrary, F is equicontinuous.

(\Leftarrow) Suppose F is bounded and equicontinuous. Let $\varepsilon > 0$. By equicontinuity, for every $x \in K$, there is some neighbourhood U_x of x such that $|f(y) - f(x)| < \varepsilon$ for all $y \in U_x$, $f \in F$. Obviously, we have

$$\bigcup_{x \in K} U_x = K.$$

By the compactness of K , there are some $\{x_1, \dots, x_n\}$ such that

$$\bigcup_{i=1}^n U_{x_i} \supseteq K.$$

Consider the restriction of functions in F to these points. This can be viewed as a bounded subset of ℓ_{∞}^n , the n -dimensional normed vector space with the supremum norm. Since this is finite-dimensional, boundedness implies total boundedness (due to, say, the compactness of the closed unit ball). In other

words, there is a finite ε -net $\{f_1, \dots, f_n\}$ such that for every $f \in F$, there is a $j \in \{1, \dots, n\}$ such that

$$\max_i |f(x_i) - f_j(x_i)| < \varepsilon.$$

Then for every $f \in F$, pick an f_j such that the above holds. Then

$$\|f - f_j\|_{C(K)} = \sup_y |f(y) - f_j(y)|$$

Since $\{U_{x_i}\}$ covers K , we can write this as

$$\begin{aligned} &= \max_i \sup_{y \in U_{x_i}} |f(y) - f_j(y)| \\ &\leq \max_i \sup_{y \in U_{x_i}} (|f(y) - f(x_i)| + |f(x_i) - f_j(x_i)| + |f_j(x_i) - f_j(y)|) \\ &< \varepsilon + \varepsilon + \varepsilon = 3\varepsilon. \end{aligned}$$

So done. □

Now we return to prove the proposition we just used to prove Arzelà-Ascoli.

Proposition. Let X be a (complete) metric space. Then $E \subseteq X$ is totally bounded if and only if for every sequence $\{y_i\}_{i=1}^\infty \subseteq E$, there is a subsequence which is Cauchy.

Proof. (\Rightarrow) Let $E \subseteq X$ be totally bounded, $\{y_i\} \in E$. For every $j \in \mathbb{N}$, there exists a finite $\frac{1}{j}$ -net, call it N_j .

Now since N_1 is finite, there is some x_1 such that there are infinitely many y_i 's in $B(x_1, 1)$. Pick the first y_i in $B(x_1, 1)$ and call it y_{i_1} .

Now there is some $x_2 \in N_2$ such that there are infinitely many y_i 's in $B(x_1, 1) \cap B(x_2, \frac{1}{2})$. Pick the one with smallest value of $i > i_1$, and call this y_{i_2} . Continue till infinity.

This procedure gives a sequence $x_i \in N_i$ and a subsequence $\{y_{i_k}\}$, and also

$$y_{i_n} \in \bigcap_{j=1}^n B\left(x_j, \frac{1}{j}\right).$$


It is easy to see that $\{y_{i_n}\}$ is Cauchy since if $m > n$, then $d(y_{i_m}, y_{i_n}) < \frac{2}{n}$.

(\Leftarrow) Suppose E is not totally bounded. So there is no finite ε -net. Pick any y_1 . Pick y_2 such that $d(y_1, y_2) \geq \varepsilon$. This exists because there is no finite ε -net.

Now given y_1, \dots, y_n such that $d(y_i, y_j) \geq \varepsilon$ for all $i, j = 1, \dots, n$, $i \neq j$, we pick y_{n+1} such that $d(y_{n+1}, y_j) \geq \varepsilon$ for all $j = 1, \dots, n$. Again, this exists because there is no finite ε -net. Then clearly any subsequence of $\{y_n\}$ is not Cauchy. □

Note that the first part is similar to the proof of Bolzano-Weierstrass in \mathbb{R}^n by repeated bisection.

Recall that at the beginning of the chapter, we have seen that boundedness and equicontinuity assumptions are necessary. The compactness of K is also important. Let $X = \mathbb{R}$, which is not compact, and let $\phi \in C_c^\infty$, an infinitely differentiable function with compact support, say, the bump function.



We now let $f_n(x) = \phi(x - n)$, i.e. we shift our bump function to the right by n units. This sequence is clearly bounded and equicontinuous, but this has no convergent subsequence — f_n converges pointwise to the zero function, but the convergence is not uniform, and this is true for arbitrary subsequences as well.

We are going to look at some applications of the theorem:

Example. Let $K \subseteq \mathbb{R}$ be a compact space, $\{f_n\}_{n=1}^\infty$ be a sequence of continuously differentiable functions in $C(K)$, such that

$$\sup_x \sup_n (|f_n(x)| + |f'_n(x)|) < C$$

for some C . Then there is a convergent subsequence. We clearly have uniform boundedness. To obtain equicontinuity, since the derivative is bounded, by the mean value theorem, we have

$$\frac{|f_n(x) - f_n(y)|}{|x - y|} = |f'_n(z)| \leq C$$

So

$$|f_n(x) - f_n(y)| \leq C|x - y|.$$

Consider the ordinary differential equation $x' = f(x)$ with the boundary conditions $x(0) = x_0 \in \mathbb{R}^n$. Recall from IB Analysis II that Picard-Lindelöf theorem says that if f is a Lipschitz function, then there exists some $\varepsilon > 0$ such that the ODE has a *unique* solution in $(-\varepsilon, \varepsilon)$.

What if f is not Lipschitz? If so, we can get the following

Theorem (Peano*). Given f continuous, then there is some $\varepsilon > 0$ such that $x' = f(x)$ with boundary condition $x(0) = x_0 \in \mathbb{R}$ has a solution in $(-\varepsilon, \varepsilon)$.

Note that uniqueness is false. For example, if $x' = \sqrt{|x|}$, $x(0) = 0$, then $x(t) = 0$ and $x(t) = t^2$ are both solutions.

Proof. (sketch) We approximate f by a sequence of continuously differentiable functions f_n such that $\|f - f_n\|_{C(K)} \rightarrow 0$ for some $K \subseteq \mathbb{R}$. We use Picard-Lindelöf to get a solution for all n . Then we use the ODE to get estimates for the solution. Finally, we can use Arzelà-Ascoli to extract a limit as $n \rightarrow \infty$. We can then show it is indeed a solution. \square

3.4 Stone–Weierstrass theorem

Here we will prove the Stone–Weierstrass theorem, which is a generalization of the classical Weierstrass approximation theorem.

Theorem (Weierstrass approximation theorem). The set of polynomials are dense in $C([0, 1])$.

This tells us that $C([0, 1])$ is not too “big” because it has a dense subset of “nice” functions.

We will try to generalize this for more general domains. Note that for this section, real-valued and complex-valued continuous functions are somewhat differentiable. So we will write these as $C_{\mathbb{R}}(K)$ and $C_{\mathbb{C}}(K)$.

To state this theorem, we need some definitions.

Definition (Algebra). A vector space $(V, +)$ is called an *algebra* if there is an operation (called multiplication) $\cdot : V \rightarrow V$ such that $(V, +, \cdot)$ is a *rng* (i.e. ring not necessarily with multiplicative identity). Also, $\lambda(\mathbf{v} \cdot \mathbf{w}) = (\lambda\mathbf{v}) \cdot \mathbf{w} = \mathbf{v} \cdot (\lambda\mathbf{w})$ for all $\lambda \in \mathbb{F}$, $\mathbf{v}, \mathbf{w} \in V$.

If V is in addition a normed vector space and

$$\|\mathbf{v} \cdot \mathbf{w}\|_V \leq \|\mathbf{v}\|_V \cdot \|\mathbf{w}\|_V$$

for all $\mathbf{v}, \mathbf{w} \in V$, then we say V is a *normed algebra*.

If V complete normed algebra, we say V is a *Banach algebra*.

If V is an algebra that is commutative as a rng, then we say V is a *commutative algebra*.

If V is an algebra with multiplicative identity, then V is a *unital algebra*.

Example. $C(K)$ is a commutative, unital, Banach algebra.

Example. Recall from the example sheets that if $T_1, T_2 : V \rightarrow V$, then

$$\|T_1 \circ T_2\|_{\mathcal{B}(V, V)} \leq \|T_1\|_{\mathcal{B}(V, V)} \|T_2\|_{\mathcal{B}(V, V)}.$$

So $\mathcal{B}(V, V)$ is a unital normed algebra.

We will need this language to generalize the Weierstrass approximation theorem. The main problem in doing so is that we have to figure out what we can generalize polynomials to. This is why we need these funny definitions.

Theorem (Stone-Weierstrass theorem). Let K be compact, and $\mathcal{A} \subseteq C_{\mathbb{R}}(K)$ be a subalgebra (i.e. it is a subset that is closed under the operations) with the property that it separates points, i.e. for every $x, y \in K$ distinct, there exists some $f \in \mathcal{A}$ such that $f(x) \neq f(y)$. Then either $\mathcal{A} = C_{\mathbb{R}}(K)$ or there is some $x_0 \in K$ such that

$$\bar{\mathcal{A}} = \{f \in C_{\mathbb{R}}(K) : f(x_0) = 0\}.$$

Note that it is not possible that $\bar{\mathcal{A}}$ is always zero on 2 or more points, since \mathcal{A} separates points.

This indeed implies the Weierstrass approximation theorem, since polynomials separates points (consider the polynomial x), and the polynomial 1 is never 0 for all x . In fact, this works also for polynomials on compact subsets of \mathbb{R}^n .

Note, however, that the second case of Stone-Weierstrass theorem can actually happen. For example, consider $K = [0, 1]$ compact, and \mathcal{A} be the algebra generated by x . Then $\bar{\mathcal{A}} = \{f \in C_{\mathbb{R}}(K) : f(0) = 0\}$.

We will prove this in using two lemmas:

Lemma. Let K compact, $\mathcal{L} \subseteq C_{\mathbb{R}}(K)$ be a subset which is closed under taking maximum and minimum, i.e. if $f, g \in \mathcal{L}$, then $\max\{f, g\} \in \mathcal{L}$ and $\min\{f, g\} \in \mathcal{L}$

(with $\max\{f, g\}$ defined as $\max\{f, g\}(x) = \max\{f(x), g(x)\}$, and similarly for minimum).

Given $g \in C_{\mathbb{R}}(K)$, assume further that for any $\varepsilon > 0$ and $x, y \in K$, there exists $f_{x,y} \in \mathcal{L}$ such that

$$|f_{x,y}(x) - g(x)| < \varepsilon, \quad |f_{x,y}(y) - g(y)| < \varepsilon.$$

Then there exists some $f \in \mathcal{L}$ such that

$$\|f - g\|_{C_{\mathbb{R}}(K)} < \varepsilon,$$

i.e. $g \in \bar{\mathcal{L}}$.

This means that if we are allowed to take maximums and minimums, to be able to approximate a function, we just need to be able to approximate it at any two points.

The idea is to next show that if \mathcal{A} is a subalgebra, then $\bar{\mathcal{A}}$ is closed under taking maximum and minimum. Then use the ability to separate points to find $f_{x,y}$, and prove that we can approximate arbitrary functions.

Proof. Let $g \in C_{\mathbb{R}}(K)$ and $\varepsilon > 0$ be given. So for every $x, y \in K$, there is some $f_{x,y} \in \mathcal{L}$ such that

$$|f_{x,y}(x) - g(x)| < \varepsilon, \quad |f_{x,y}(y) - g(y)| < \varepsilon.$$

Claim. For each $x \in K$, there exists $f_x \in \mathcal{L}$ such that $|f_x(x) - g(x)| < \varepsilon$ and $f_x(z) < g(z) + \varepsilon$ for all $z \in K$.

Since $f_{x,y}$ is continuous, there is some $U_{x,y}$ containing y open such that


$$|f_{x,y}(z) - g(z)| < \varepsilon$$

for all $z \in U_{x,y}$. Since

$$\bigcup_{y \in K} U_{x,y} \supseteq K,$$

by compactness of K , there exists a some y_1, \dots, y_n such that

$$\bigcup_{i=1}^n U_{x,y_i} \supseteq K.$$



We then let

$$f_x(z) = \min\{f_{x,y_1}(z), \dots, f_{x,y_n}(z)\}$$

for every $z \in K$. We then see that this works. Indeed, by assumption, $f_x \in \mathcal{L}$. If $z \in K$ is some arbitrary point, then $z \in U_{x,y_i}$ for some i . Then

$$f_{x,y_i}(z) < g(z) + \varepsilon.$$

Hence, since f_x is the minimum of all such f_{x,y_i} , for any z , we have

$$f_x(z) < g(z) + \varepsilon.$$

The property at x is also clear.

Claim. There exists $f \in \mathcal{L}$ such that $|f(z) - g(z)| < \varepsilon$ for all $z \in K$.

We are going to play the same game with this. By continuity of f_x , there is V_x containing x open such that

$$|f_x(w) - g(w)| < \varepsilon$$

for all $w \in V_x$. Since

$$\bigcup_{x \in K} V_x \supseteq K,$$

by compactness of K , there is some $\{x_1, \dots, x_m\}$ such that

$$\bigcup_{j=1}^m V_{x_j} \supseteq K.$$

Define

$$f(z) = \max\{f_{x_1}(z), \dots, f_{x_m}(z)\}.$$

Again, by assumption, $f \in \mathcal{L}$. Then we know that

$$f(z) > g(z) - \varepsilon.$$

We still have our first bound

$$f(z) < g(z) + \varepsilon.$$

Therefore we have

$$\|f - g\|_{C_{\mathbb{R}}(K)} < \varepsilon. \quad \square$$

Lemma. Let $\mathcal{A} \subseteq C_{\mathbb{R}}(K)$ be a subalgebra that is a closed subset in the topology of $C_{\mathbb{R}}(K)$. Then \mathcal{A} is closed under taking maximum and minimum.

Proof. First note that

$$\begin{aligned} \max\{f(x), g(x)\} &= \frac{1}{2}(f(x) + g(x)) + \frac{1}{2}|f(x) - g(x)|, \\ \min\{f(x), g(x)\} &= \frac{1}{2}(f(x) + g(x)) - \frac{1}{2}|f(x) - g(x)|. \end{aligned}$$

Since \mathcal{A} is an algebra, it suffices to show that $f \in \mathcal{A}$ implies $|f| \in \mathcal{A}$ for every f such that $\|f\|_{C_{\mathbb{R}}(K)} \leq 1$.

The key observation is the following: consider the function $h(x) = \sqrt{x + \varepsilon^2}$. Then $h(x^2)$ approximates $|x|$. This has the property that the Taylor expansion of $h(x)$ centered at $x = \frac{1}{2}$ is uniformly convergent for $x \in [0, 1]$. Therefore there exists a polynomial $S(x)$ such that

$$|S(x) - \sqrt{x + \varepsilon^2}| < \varepsilon.$$

Now note that $S(x) - S(0)$ is a polynomial with no constant term. Therefore, since \mathcal{A} is an algebra, if $f \in \mathcal{A}$, then $S(f^2) - S(0) \in \mathcal{A}$ by closure.

Now look at

$$\| |f| - (S(f^2) - S(0)) \|_{C_{\mathbb{R}}(K)} \leq \| |f| - \sqrt{f^2 + \varepsilon^2} \| + \| \sqrt{f^2 + \varepsilon^2} - S(f^2) \| + \| S(0) \|.$$

We will make each individual term small. For the first term, note that

$$\sup_{x \in [0,1]} |x - \sqrt{x^2 + \varepsilon^2}| = \sup_{x \in [0,1]} \frac{\varepsilon^2}{|x + \sqrt{x^2 + \varepsilon^2}|} = \varepsilon.$$

So the first term is at most ε . The second term is also easy, since S is chosen such that $|S(x) - \sqrt{x + \varepsilon^2}| < 1$ for $x \in [0, 1]$, and $|f(x)^2| \leq 1$ for all $x \in [0, 1]$. So it is again bounded by ε .

By the same formula, $|S(0) - \sqrt{0 + \varepsilon^2}| < \varepsilon$. So $|S(0)| < 2\varepsilon$. So

$$\| |f| - (S(f^2) - S(0)) \|_{C_{\mathbb{R}}(K)} < 4\varepsilon.$$

Since $\varepsilon > 0$ and \mathcal{A} is closed in the topology of $C_{\mathbb{R}}(K)$, $f \in \mathcal{A}$ and $\|f\|_{C_{\mathbb{R}}(K)} \leq 1$ implies that $|f| \in \mathcal{A}$. \square

Note that if we have already proven the classical Weierstrass approximation theorem, we can just use it to get a polynomial approximation for $|f|$ directly.

We will now combine both lemmas and prove the Stone-Weierstrass theorem.

Theorem (Stone-Weierstrass theorem). Let K be compact, and $\mathcal{A} \subseteq C_{\mathbb{R}}(K)$ be a subalgebra (i.e. it is a subset that is closed under the operations) with the property that it separates points, i.e. for every $x, y \in K$ distinct, there exists some $f \in \mathcal{A}$ such that $f(x) \neq f(y)$. Then either $\bar{\mathcal{A}} = C_{\mathbb{R}}(K)$ or there is some $x_0 \in K$ such that

$$\bar{\mathcal{A}} = \{f \in C_{\mathbb{R}}(K) : f(x_0) = 0\}.$$

Proof. Note that there are two possible outcomes. We will first look at the first possibility.

Consider the case where for all $x \in K$, there is some $f \in \mathcal{A}$ such that $f(x) \neq 0$. Let $g \in C_{\mathbb{R}}(K)$ be given. By our previous lemmas, to approximate g in $\bar{\mathcal{A}}$, we just need to show that we can approximate g at two points. So given any $\varepsilon > 0$, $x, y \in K$, we want to find $f_{x,y} \in \mathcal{A}$ such that

$$|f_{x,y}(x) - g(x)| < \varepsilon, \quad |f_{x,y}(y) - g(y)| < \varepsilon. \quad (*)$$

For every $x, y \in K$, $x \neq y$, we first show that there exists $h_{x,y} \in \mathcal{A}$ such that $h_{x,y}(x) \neq 0$, and $h_{x,y}(x) \neq h_{x,y}(y)$. This is easy to see. By our assumptions, we can find the following functions:

- (i) There exists $h_{x,y}^{(1)}$ such that $h_{x,y}^{(1)} \neq h_{x,y}^{(1)}(y)$

(ii) There exists $h_{x,y}^{(2)}$ such that $h_{x,y}^{(2)}(x) \neq 0$.

(iii) There exists $h_{x,y}^{(3)}$ such that $h_{x,y}^{(3)}(y) \neq 0$.

Then it is an easy exercise to show that some linear combination of $h_{x,y}^{(1)}$ and $h_{x,y}^{(2)}$ and $h_{x,y}^{(3)}$ works, say $h_{x,y}$.

We will want to find our $f_{x,y}$ that satisfies (*). But we will do better. We will make it equal g on x and y . The idea is to take linear combinations of $h_{x,y}$ and $h_{x,y}^2$. Instead of doing the messy algebra to show that we can find a working linear combination, just notice that $(h_{x,y}(x), h_{x,y}(y))$ and $(h_{x,y}(x)^2, h_{x,y}(y)^2)$ are linearly independent vectors in \mathbb{R}^2 . Therefore there exists $\alpha, \beta \in \mathbb{R}$ such that

$$\alpha(h_{x,y}(x), h_{x,y}(y)) + \beta(h_{x,y}(x)^2, h_{x,y}(y)^2) = (g(x), g(y)).$$

So done.

In the other case, given \mathcal{A} , suppose there is $x_0 \in K$ such that $f(x_0) = 0$ for all $f \in \mathcal{A}$. Consider the algebra

$$\mathcal{A}' = \mathcal{A} + \lambda 1 = \{f + \lambda 1 : f \in \mathcal{A}, \lambda \in \mathbb{R}\}$$

Since \mathcal{A} separates points, and for any $x \in K$, there is some $f \in \mathcal{A}'$ such that $f(x) \neq 0$ (e.g. $f = 1$), by the previous part, we know that $\bar{\mathcal{A}}' = C_{\mathbb{R}}(K)$.

Now note that

$$\bar{\mathcal{A}} \subseteq \{f \in C_{\mathbb{R}}(K) : f(x_0) = 0\} = B.$$

So we suffice to show that we have equality, i.e. for any $g \in B$ and $\varepsilon > 0$, there is some $f \in \mathcal{A}$ such that

$$\|f - g\|_{C_{\mathbb{R}}(K)} < \varepsilon.$$

Since $\bar{\mathcal{A}}' = C_{\mathbb{R}}(K)$, given such g and ε , there is some $f \in \mathcal{A}$ and $\lambda_0 \in \mathbb{R}$ such that

$$\|g - (f + \lambda 1)\|_{C_{\mathbb{R}}(K)} < \varepsilon.$$

But $g(x_0) = f(x_0) = 0$, which implies that $|\lambda| < \varepsilon$. Therefore $\|g - f\|_{C_{\mathbb{R}}(K)} < 2\varepsilon$. So done. \square

What happens for $C_{\mathbb{C}}(K)$?

Example. Let $K = \overline{B(0, 1)} \subseteq \mathbb{C}$, and let \mathcal{A} be the set of polynomials on $\overline{B(0, 1)}$. We will show that $\bar{\mathcal{A}} \neq C_{\mathbb{C}}(\overline{B(0, 1)})$.

Consider $f(z) = \bar{z}$. This is not in the closure of \mathcal{A} , since this is not holomorphic, but the uniform limit of any sequence of holomorphic is holomorphic (by Morera's theorem, i.e. f is holomorphic iff $\oint_{\gamma} f(z) dz = 0$ for all closed piecewise smooth curve γ).

Hence we need an additional condition on this. It turns out that this rather simple example is the only way in which things can break. We have the following:

Theorem (Complex version of Stone-Weierstrass theorem). Let K be compact and $\mathcal{A} \subseteq C_{\mathbb{C}}(K)$ be a subalgebra over \mathbb{C} which separates points and is closed under complex conjugation (i.e. if $f \in \mathcal{A}$, then $\bar{f} \in \mathcal{A}$). Then either $\bar{\mathcal{A}} = C_{\mathbb{C}}(K)$ or there is an x_0 such that $\bar{\mathcal{A}} = \{f \in C_{\mathbb{C}}(K) : f(x_0) = 0\}$.

Proof. It suffices to show that either $\bar{\mathcal{A}} \supseteq C_{\mathbb{R}}(K)$ or there exists a point x_0 such that $\bar{\mathcal{A}} \supseteq \{f \in C_{\mathbb{R}}(K) : f(x_0) = 0\}$, since we can always break a complex function up into its real and imaginary parts.

Now consider

$$\mathcal{A}' = \left\{ \frac{f + \bar{f}}{2} : f \in \mathcal{A} \right\} \cup \left\{ \frac{f - \bar{f}}{2i} : f \in \mathcal{A} \right\}.$$

Now note that by closure of \mathcal{A} , we know that \mathcal{A}' is a subset of \mathcal{A} and is a subalgebra of $C_{\mathbb{R}}(K)$ over \mathbb{R} , which separates points. Hence by the real version of Stone-Weierstrass, either $\bar{\mathcal{A}}' = C_{\mathbb{R}}(K)$ or there is some x_0 such that $\bar{\mathcal{A}}' = \{f \in C_{\mathbb{R}}(K) : f(x_0) = 0\}$. So done. \square

4 Hilbert spaces

4.1 Inner product spaces

We have just looked at continuous functions on some compact space K . Another important space is the space of square-integrable functions. Consider the space

$$L^2(\mathbb{R}) = \left\{ f : f \text{ is Lebesgue integrable} : \int |f|^2 < \infty \right\} / \sim,$$

where $f \sim g$ if $f = g$ is Lebesgue almost everywhere.

One thing we like to think about is the Fourier series. Recall that for $f \in C(S^1)$, we have defined, for each $k \in \mathbb{Z}$,

$$\hat{f}(k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ikx} f(x) dx,$$

and we have defined the partial sum

$$S_N(f)(x) = \sum_{n=-N}^N e^{inx} \hat{f}(k).$$

We have previously seen that even if f is continuous, it is possible that the partial sums S_N do not converge, even pointwise. However, we can ask for something weaker:

Proposition. Let $f \in C(S^1)$. Then

$$\lim_{N \rightarrow \infty} \frac{1}{2\pi} \int_{-\pi}^{\pi} |f(x) - S_N(f)(x)|^2 dx = 0.$$

We will prove this later. However, the key points of the proof is the “orthogonality” of $\{e^{inx}\}$. More precisely, we have

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{inx} e^{-imx} dx = 0 \text{ if } m \neq n.$$

The introduction of Hilbert spaces is in particular a way to put this in a general framework. We want to introduce an extra structure that gives rise to “orthogonality”.

Definition (Inner product). Let V be a vector space over \mathbb{R} or \mathbb{C} . We say $p : V \times V \rightarrow \mathbb{R}$ or \mathbb{C} is an *inner product* on V it satisfies

- (i) $p(\mathbf{v}, \mathbf{w}) = \overline{p(\mathbf{w}, \mathbf{v})}$ for all $\mathbf{v}, \mathbf{w} \in V$. (antisymmetry)
- (ii) $p(\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2, \mathbf{u}) = \lambda_1 p(\mathbf{v}_1, \mathbf{u}) + \lambda_2 p(\mathbf{v}_2, \mathbf{u})$. (linearity in first argument)
- (iii) $p(\mathbf{v}, \mathbf{v}) \geq 0$ for all $\mathbf{v} \in V$ and equality holds iff $\mathbf{v} = \mathbf{0}$. (non-negativity)

We will often denote an inner product by $p(\mathbf{v}, \mathbf{w}) = \langle \mathbf{v}, \mathbf{w} \rangle$. We call $(V, \langle \cdot, \cdot \rangle)$ an *inner product space*.

Definition (Orthogonality). In an inner product space, \mathbf{v} and \mathbf{w} are *orthogonal* if $\langle \mathbf{v}, \mathbf{w} \rangle = 0$.

Orthogonality and the inner product is important when dealing with vector spaces. For example, recall that when working with finite-dimensional spaces, we had things like Hermitian matrices, orthogonal matrices and normal matrices. All these are in some sense defined in terms of the inner product and orthogonality. More fundamentally, when we have a finite-dimensional vector spaces, we often write the vectors as a set of n coordinates. To define this coordinate system, we start by picking n orthogonal vectors (which are in fact orthonormal), and then the coordinates are just the projections onto these orthogonal vectors.

Hopefully, you are convinced that inner products are important. So let's see what we can get if we put in inner products to arbitrary vector spaces.

We will look at some easy properties of the inner product.

Proposition (Cauchy-Schwarz inequality). Let $(V, \langle \cdot, \cdot \rangle)$ be an inner product space. Then for all $\mathbf{v}, \mathbf{w} \in V$,

$$|\langle \mathbf{v}, \mathbf{w} \rangle| \leq \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle \langle \mathbf{w}, \mathbf{w} \rangle},$$

with equality iff there is some $\lambda \in \mathbb{R}$ or \mathbb{C} such that $\mathbf{v} = \lambda \mathbf{w}$ or $\mathbf{w} = \lambda \mathbf{v}$.

Proof. wlog, we can assume $\mathbf{w} \neq 0$. Otherwise, this is trivial. Moreover, assume $\langle \mathbf{v}, \mathbf{w} \rangle \in \mathbb{R}$. Otherwise, we can just multiply \mathbf{w} by some $e^{i\alpha}$.

By non-negativity, we know that for all t , we have

$$\begin{aligned} 0 &\leq \langle \mathbf{v} + t\mathbf{w}, \mathbf{v} + t\mathbf{w} \rangle \\ &= \langle \mathbf{v}, \mathbf{v} \rangle + 2t\langle \mathbf{v}, \mathbf{w} \rangle + t^2\langle \mathbf{w}, \mathbf{w} \rangle. \end{aligned}$$

Therefore, the discriminant of this quadratic polynomial in t is non-positive, i.e.

$$4(\langle \mathbf{v}, \mathbf{w} \rangle)^2 - 4\langle \mathbf{v}, \mathbf{v} \rangle \langle \mathbf{w}, \mathbf{w} \rangle \leq 0,$$

from which the result follows.

Finally, note that if equality holds, then the discriminant is 0. So the quadratic has exactly one root. So there exists t such that $\mathbf{v} + t\mathbf{w} = 0$, which of course implies $\mathbf{v} = -t\mathbf{w}$. \square

Proposition. Let $(V, \langle \cdot, \cdot \rangle)$ be an inner product space. Then

$$\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$$

defines a norm.

Proof. The first two axioms of the norm are easy to check, since it follows directly from definition of the inner product that $\|\mathbf{v}\| \geq 0$ with equality iff $\mathbf{v} = \mathbf{0}$, and $\|\lambda \mathbf{v}\| = |\lambda| \|\mathbf{v}\|$.

The only non-trivial thing to check is the triangle inequality. We have

$$\begin{aligned} \|\mathbf{v} + \mathbf{w}\|^2 &= \langle \mathbf{v} + \mathbf{w}, \mathbf{v} + \mathbf{w} \rangle \\ &= \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 + |\langle \mathbf{v}, \mathbf{w} \rangle| + |\langle \mathbf{w}, \mathbf{v} \rangle| \\ &\leq \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 + 2\|\mathbf{v}\|\|\mathbf{w}\| \\ &= (\|\mathbf{v}\| + \|\mathbf{w}\|)^2 \end{aligned}$$

Hence we know that $\|\mathbf{v} + \mathbf{w}\| \leq \|\mathbf{v}\| + \|\mathbf{w}\|$. \square

This motivates the following definition:

Definition (Euclidean space). A normed vector space $(V, \|\cdot\|)$ is a *Euclidean space* if there exists an inner product $\langle \cdot, \cdot \rangle$ such that

$$\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}.$$

Proposition. Let $(E, \|\cdot\|)$ be a Euclidean space. Then there is a *unique* inner product $\langle \cdot, \cdot \rangle$ such that $\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$.

Proof. The real and complex cases are slightly different.

First suppose E is a vector space over \mathbb{R} , and suppose also that we have an inner product $\langle \cdot, \cdot \rangle$ such that $\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$. Then

$$\langle \mathbf{v} + \mathbf{w}, \mathbf{v} + \mathbf{w} \rangle = \|\mathbf{v}\|^2 + 2\langle \mathbf{v}, \mathbf{w} \rangle + \|\mathbf{w}\|^2.$$

So we get

$$\langle \mathbf{v}, \mathbf{w} \rangle = \frac{1}{2}(\|\mathbf{v} + \mathbf{w}\|^2 - \|\mathbf{v}\|^2 - \|\mathbf{w}\|^2). \quad (*)$$

In particular, the inner product is completely determined by the norm. So this must be unique.

Now suppose E is a vector space over \mathbb{C} . We have

$$\langle \mathbf{v} + \mathbf{w}, \mathbf{v} + \mathbf{w} \rangle = \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 + \langle \mathbf{v}, \mathbf{w} \rangle + \langle \mathbf{w}, \mathbf{v} \rangle \quad (1)$$

$$\langle \mathbf{v} - \mathbf{w}, \mathbf{v} - \mathbf{w} \rangle = \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 - \langle \mathbf{v}, \mathbf{w} \rangle - \langle \mathbf{w}, \mathbf{v} \rangle \quad (2)$$

$$\langle \mathbf{v} + i\mathbf{w}, \mathbf{v} + i\mathbf{w} \rangle = \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 - i\langle \mathbf{v}, \mathbf{w} \rangle + i\langle \mathbf{w}, \mathbf{v} \rangle \quad (3)$$

$$\langle \mathbf{v} - i\mathbf{w}, \mathbf{v} - i\mathbf{w} \rangle = \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 + i\langle \mathbf{v}, \mathbf{w} \rangle - i\langle \mathbf{w}, \mathbf{v} \rangle \quad (4)$$

Now consider $(1) - (2) + i(3) - i(4)$. Then we obtain

$$\|\mathbf{v} + \mathbf{w}\|^2 - \|\mathbf{v} - \mathbf{w}\|^2 + i\|\mathbf{v} + i\mathbf{w}\|^2 - i\|\mathbf{v} - i\mathbf{w}\|^2 = 4\langle \mathbf{v}, \mathbf{w} \rangle. \quad (\dagger)$$

So again $\langle \mathbf{v}, \mathbf{w} \rangle$ is again determined by the norm. \square

The identities $(*)$ and (\dagger) are sometimes known as the *polarization identities*.


Definition (Hilbert space). A Euclidean space $(E, \|\cdot\|)$ is a *Hilbert space* if it is complete.

We will prove certain properties of the inner product.

Proposition (Parallelogram law). Let $(E, \|\cdot\|)$ be a Euclidean space. Then for $\mathbf{v}, \mathbf{w} \in E$, we have

$$\|\mathbf{v} - \mathbf{w}\|^2 + \|\mathbf{v} + \mathbf{w}\|^2 = 2\|\mathbf{v}\|^2 + 2\|\mathbf{w}\|^2.$$

This is called the parallelogram law because it says that for any parallelogram, the sum of the square of the lengths of the diagonals is the sum of square of the lengths of the two sides.



Proof. This is just simple algebraic manipulation. We have

$$\begin{aligned}\|\mathbf{v} - \mathbf{w}\|^2 + \|\mathbf{v} + \mathbf{w}\|^2 &= \langle \mathbf{v} - \mathbf{w}, \mathbf{v} - \mathbf{w} \rangle + \langle \mathbf{v} + \mathbf{w}, \mathbf{v} + \mathbf{w} \rangle \\ &= \langle \mathbf{v}, \mathbf{v} \rangle - \langle \mathbf{v}, \mathbf{w} \rangle - \langle \mathbf{w}, \mathbf{v} \rangle + \langle \mathbf{w}, \mathbf{w} \rangle \\ &\quad + \langle \mathbf{v}, \mathbf{v} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle + \langle \mathbf{w}, \mathbf{v} \rangle + \langle \mathbf{w}, \mathbf{w} \rangle \\ &= 2\langle \mathbf{v}, \mathbf{v} \rangle + 2\langle \mathbf{w}, \mathbf{w} \rangle.\end{aligned}\quad \square$$

Proposition (Pythagoras theorem). Let $(E, \|\cdot\|)$ be a Euclidean space, and let $\mathbf{v}, \mathbf{w} \in E$ be orthogonal. Then

$$\|\mathbf{v} + \mathbf{w}\|^2 = \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2.$$

Proof.

$$\begin{aligned}\|\mathbf{v} + \mathbf{w}\|^2 &= \langle \mathbf{v} + \mathbf{w}, \mathbf{v} + \mathbf{w} \rangle \\ &= \langle \mathbf{v}, \mathbf{v} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle + \langle \mathbf{w}, \mathbf{v} \rangle + \langle \mathbf{w}, \mathbf{w} \rangle \\ &= \langle \mathbf{v}, \mathbf{v} \rangle + 0 + 0 + \langle \mathbf{w}, \mathbf{w} \rangle \\ &= \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2.\end{aligned}\quad \square$$

By induction, if $\mathbf{v}_i \in E$ for $i = 1, \dots, n$ such that $\langle \mathbf{v}_i, \mathbf{v}_j \rangle = 0$ for $i \neq j$, i.e. they are mutually orthogonal, then

$$\left\| \sum_{i=1}^n \mathbf{v}_i \right\|^2 = \sum_{i=1}^n \|\mathbf{v}_i\|^2.$$

Proposition. Let $(E, \|\cdot\|)$ be a Euclidean space. Then $\langle \cdot, \cdot \rangle : E \times E \rightarrow \mathbb{C}$ is continuous.

Proof. Let $(\mathbf{v}, \mathbf{w}) \in E \times E$, and $(\tilde{\mathbf{v}}, \tilde{\mathbf{w}}) \in E \times E$. We have

$$\begin{aligned}\|\langle \mathbf{v}, \mathbf{w} \rangle - \langle \tilde{\mathbf{v}}, \tilde{\mathbf{w}} \rangle\| &= \|\langle \mathbf{v}, \mathbf{w} \rangle - \langle \mathbf{v}, \tilde{\mathbf{w}} \rangle + \langle \mathbf{v}, \tilde{\mathbf{w}} \rangle - \langle \tilde{\mathbf{v}}, \tilde{\mathbf{w}} \rangle\| \\ &\leq \|\langle \mathbf{v}, \mathbf{w} \rangle - \langle \mathbf{v}, \tilde{\mathbf{w}} \rangle\| + \|\langle \mathbf{v}, \tilde{\mathbf{w}} \rangle - \langle \tilde{\mathbf{v}}, \tilde{\mathbf{w}} \rangle\| \\ &= \|\langle \mathbf{v}, \mathbf{w} - \tilde{\mathbf{w}} \rangle\| + \|\langle \mathbf{v} - \tilde{\mathbf{v}}, \tilde{\mathbf{w}} \rangle\| \\ &\leq \|\mathbf{v}\| \|\mathbf{w} - \tilde{\mathbf{w}}\| + \|\mathbf{v} - \tilde{\mathbf{v}}\| \|\tilde{\mathbf{w}}\|\end{aligned}$$

Hence for \mathbf{v}, \mathbf{w} sufficiently closed to $\tilde{\mathbf{v}}, \tilde{\mathbf{w}}$, we can get $\|\langle \mathbf{v}, \mathbf{w} \rangle - \langle \tilde{\mathbf{v}}, \tilde{\mathbf{w}} \rangle\|$ arbitrarily small. So it is continuous. \square

When we have an incomplete Euclidean space, we can of course take the completion of it to form a complete extension of the original normed vector space. However, it is not immediately obvious that the inner product can also be extended to the completion to give a Hilbert space. The following proposition tells us we can do so.

Proposition. Let $(E, \|\cdot\|)$ denote a Euclidean space, and \bar{E} its completion. Then the inner product extends to an inner product on \bar{E} , turning \bar{E} into a Hilbert space.

Proof. Recall we constructed the completion of a space as the equivalence classes of Cauchy sequences (where two Cauchy sequences (x_n) and (x'_n) are equivalent if $|x_n - x'_n| \rightarrow 0$). Let $(\mathbf{x}_n), (\mathbf{y}_n)$ be two Cauchy sequences in E , and let $\tilde{x}, \tilde{y} \in \bar{E}$ denote their equivalence classes. We define the inner product as

$$\langle \tilde{\mathbf{x}}, \tilde{\mathbf{y}} \rangle = \lim_{n \rightarrow \infty} \langle \mathbf{x}_n, \mathbf{y}_n \rangle. \quad (*)$$

We want to show this is well-defined. Firstly, we need to make sure the limit exists. We can show this by showing that this is a Cauchy sequence. We have

$$\begin{aligned} \|\langle \mathbf{x}_n, \mathbf{y}_n \rangle - \langle \mathbf{x}_m, \mathbf{y}_m \rangle\| &= \|\langle \mathbf{x}_n, \mathbf{y}_n \rangle - \langle \mathbf{x}_m, \mathbf{y}_n \rangle + \langle \mathbf{x}_m, \mathbf{y}_n \rangle - \langle \mathbf{x}_m, \mathbf{y}_m \rangle\| \\ &\leq \|\langle \mathbf{x}_n, \mathbf{y}_n \rangle - \langle \mathbf{x}_m, \mathbf{y}_n \rangle\| + \|\langle \mathbf{x}_m, \mathbf{y}_n \rangle - \langle \mathbf{x}_m, \mathbf{y}_m \rangle\| \\ &\leq \|\langle \mathbf{x}_n, \mathbf{x}_m, \mathbf{y}_n \rangle\| + \|\langle \mathbf{x}_m, \mathbf{y}_n - \mathbf{y}_m \rangle\| \\ &\leq \|\mathbf{x}_n - \mathbf{x}_m\| \|\mathbf{y}_n\| + \|\mathbf{x}\| \|\mathbf{y}_n - \mathbf{y}_m\| \end{aligned}$$

So $\langle \mathbf{x}_n, \mathbf{y}_n \rangle$ is a Cauchy sequence since (\mathbf{x}_n) and (\mathbf{y}_n) are.

We also need to show that $(*)$ does not depend on the representatives for $\tilde{\mathbf{x}}$ and $\tilde{\mathbf{y}}$. This is left as an exercise for the reader

We also need to show that $\langle \cdot, \cdot \rangle_{\bar{E}}$ define the norm of $\|\cdot\|_{\bar{E}}$, which is yet another exercise. \square

Example. Consider the space

$$\ell_2 = \left\{ (x_1, x_2, \dots) : x_i \in \mathbb{C}, \sum_{i=1}^{\infty} |x_i|^2 < \infty \right\}.$$

We already know that this is a complete Banach space. We can also define an inner product on this space by

$$\langle \mathbf{a}, \mathbf{b} \rangle_{\ell^2} = \sum_{i=1}^{\infty} a_i \bar{b}_i.$$

We need to check that this actually converges. We prove this by showing absolute convergence. For each n , we can use Cauchy-Schwarz to obtain

$$\sum_{i=1}^n |a_i \bar{b}_i| \leq \left(\sum_{i=1}^n |a_i|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^n |b_i|^2 \right)^{\frac{1}{2}} \leq \|\mathbf{a}\|_{\ell^2} \|\mathbf{b}\|_{\ell^2}.$$

So it converges. Now notice that the ℓ^2 norm is indeed induced by this inner product.

This is a significant example since we will later show that every separable (i.e. has countable basis) infinite dimensional Hilbert space is isometric isomorphic to ℓ^2 .

Definition (Orthogonal space). Let E be a Euclidean space and $S \subseteq E$ an arbitrary subset. Then the *orthogonal space* of S , denoted by S^\perp is given by

$$S^\perp = \{ \mathbf{v} \in E : \forall \mathbf{w} \in S, \langle \mathbf{v}, \mathbf{w} \rangle = 0 \}.$$

Proposition. Let E be a Euclidean space and $S \subseteq E$. Then S^\perp is a closed subspace of E , and moreover

$$S^\perp = (\overline{\text{span } S})^\perp.$$

Proof. We first show it is a subspace. Let $\mathbf{u}, \mathbf{v} \in S^\perp$ and $\lambda, \mu \in \mathbb{C}$. We want to show $\lambda\mathbf{u} + \mu\mathbf{v} \in S^\perp$. Let $\mathbf{w} \in S$. Then

$$\langle \lambda\mathbf{u} + \mu\mathbf{v}, \mathbf{w} \rangle = \lambda\langle \mathbf{u}, \mathbf{w} \rangle + \mu\langle \mathbf{v}, \mathbf{w} \rangle = 0.$$

To show it is closed, let $\mathbf{u}_n \in S^\perp$ be a sequence such that $\mathbf{u}_n \rightarrow \mathbf{u} \in E$. Let $\mathbf{w} \in S$. Then we know that

$$\langle \mathbf{u}_n, \mathbf{w} \rangle = 0.$$

Hence, by the continuity of the inner product, we have

$$0 = \lim_{n \rightarrow \infty} \langle \mathbf{u}_n, \mathbf{w} \rangle = \langle \lim \mathbf{u}_n, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle.$$


The remaining part is left as an exercise. □

Note that if V is a linear subspace, then $V \cap V^\perp = \{0\}$, since any $\mathbf{v} \in V \cap V^\perp$ has to satisfy $\langle \mathbf{v}, \mathbf{v} \rangle = 0$. So $V + V^\perp$ is a direct sum.

Theorem. Let $(E, \|\cdot\|)$ be a Euclidean space, and $F \subseteq E$ a *complete* subspace. Then $F \oplus F^\perp = E$.

Hence, by definition of the direct sum, for $\mathbf{x} \in E$, we can write $\mathbf{x} = \mathbf{x}_1 + \mathbf{x}_2$, where $\mathbf{x}_1 \in F$ and $\mathbf{x}_2 \in F^\perp$. Moreover, \mathbf{x}_1 is uniquely characterized by

$$\|\mathbf{x}_1 - \mathbf{x}\| = \inf_{\mathbf{y} \in F} \|\mathbf{y} - \mathbf{x}\|.$$



Note that this is not necessarily true if F is not complete.

Proof. We already know that $F \oplus F^\perp$ is a direct sum. It thus suffices to show that the sum is the whole of E .

Let $\mathbf{y}_i \in F$ be a sequence with

$$\lim_{i \rightarrow \infty} \|\mathbf{y}_i - \mathbf{x}\| = \inf_{\mathbf{y} \in F} \|\mathbf{y} - \mathbf{x}\| = d.$$

We want to show that \mathbf{y} is a Cauchy sequence. Let $\varepsilon > 0$ be given. Let $n_0 \in \mathbb{N}$ such that for all $i \geq n_0$, we have

$$\|\mathbf{y}_i - \mathbf{x}\|^2 \leq d^2 + \varepsilon.$$

We now use the parallelogram law for $\mathbf{v} = \mathbf{x} - \mathbf{y}_i$, $\mathbf{w} = \mathbf{x} - \mathbf{y}_j$ with $i, j \geq n_0$. Then the parallelogram law says:

$$\|\mathbf{v} + \mathbf{w}\|^2 + \|\mathbf{v} - \mathbf{w}\|^2 = 2\|\mathbf{v}\|^2 + 2\|\mathbf{w}\|^2,$$

or

$$\|\mathbf{y}_j - \mathbf{y}_i\|^2 + \|2\mathbf{x} - \mathbf{y}_i - \mathbf{y}_j\|^2 = 2\|\mathbf{y}_i - \mathbf{x}\|^2 + 2\|\mathbf{y}_j - \mathbf{x}\|^2.$$

Hence we know that

$$\begin{aligned} \|\mathbf{y}_i - \mathbf{y}_j\|^2 &\leq 2\|\mathbf{y}_i - \mathbf{x}\|^2 + 2\|\mathbf{y}_j - \mathbf{x}\|^2 - 4\left\|\mathbf{x} - \frac{\mathbf{y}_i + \mathbf{y}_j}{2}\right\|^2 \\ &\leq 2(d^2 + \varepsilon) + 2(d^2 + \varepsilon) - 4d^2 \\ &\leq 4\varepsilon. \end{aligned}$$

So \mathbf{y}_i is a Cauchy sequence. Since F is complete, $\mathbf{y}_i \rightarrow \mathbf{y} \in F$ for some F . Moreover, by continuity, of $\|\cdot\|$, we know that

$$d = \lim_{i \rightarrow \infty} \|\mathbf{y}_i - \mathbf{x}\| = \|\mathbf{y} - \mathbf{x}\|.$$

Now let $\mathbf{x}_1 = \mathbf{y}$ and $\mathbf{x}_2 = \mathbf{x} - \mathbf{y}$. The only thing left over is to show $\mathbf{x}_2 \in F^\perp$. Suppose not. Then there is some $\tilde{\mathbf{y}} \in F$ such that

$$\langle \tilde{\mathbf{y}}, \mathbf{x}_2 \rangle \neq 0.$$

The idea is that we can perturb \mathbf{y} by a little bit to get a point even closer to \mathbf{x} . By multiplying $\tilde{\mathbf{y}}$ with a scalar, we can assume

$$\langle \tilde{\mathbf{y}}, \mathbf{x}_2 \rangle > 0.$$

Then for $t > 0$, we have

$$\begin{aligned} \|(\mathbf{y} + t\tilde{\mathbf{y}}) - \mathbf{x}\|^2 &= \langle \mathbf{y} + t\tilde{\mathbf{y}} - \mathbf{x}, \mathbf{y} + t\tilde{\mathbf{y}} - \mathbf{x} \rangle \\ &= \langle \mathbf{y} - \mathbf{x}, \mathbf{y} - \mathbf{x} \rangle + \langle t\tilde{\mathbf{y}}, \mathbf{y} - \mathbf{x} \rangle + \langle \mathbf{y} - \mathbf{x}, t\tilde{\mathbf{y}} \rangle + t^2 \langle \tilde{\mathbf{y}}, \tilde{\mathbf{y}} \rangle \\ &= d^2 - 2t \langle \tilde{\mathbf{y}}, \mathbf{x}_2 \rangle + t^2 \|\tilde{\mathbf{y}}\|^2. \end{aligned}$$

Hence for sufficiently small t , the t^2 term is negligible, and we can make this less than d^2 . This is a contradiction since $\mathbf{y} + t\tilde{\mathbf{y}} \in F$. \square

As a corollary, we can define the *projection map* as follows:

Corollary. Let E be a Euclidean space and $F \subseteq E$ a complete subspace. Then there exists a projection map $P : E \rightarrow E$ defined by $P(\mathbf{x}) = \mathbf{x}_1$, where $\mathbf{x}_1 \in F$ is as defined in the theorem above. Moreover, P satisfies the following properties:

- (i) $P(E) = F$ and $P(F^\perp) = \{0\}$, and $P^2 = P$. In other words, $F^\perp \leq \ker P$.
- (ii) $(I - P)(E) = F^\perp$, $(I - P)(F) = \{0\}$, $(I - P)^2 = (I - P)$.
- (iii) $\|P\|_{\mathcal{B}(E,E)} \leq 1$ and $\|I - P\|_{\mathcal{B}(E,E)} \leq 1$, with equality if and only if $F \neq \{0\}$ and $F^\perp \neq \{0\}$ respectively.

Here P projects our space to F , where $I - P$ projects our space to F^\perp .

4.2 Riesz representation theorem

Our next theorem allows us to completely understand the duals of Hilbert spaces.

Consider the following map. Given a Hilbert space H and $\mathbf{v} \in H$, consider $\phi_{\mathbf{v}} \in H^*$ defined by

$$\phi_{\mathbf{v}}(\mathbf{w}) = \langle \mathbf{w}, \mathbf{v} \rangle.$$

Note that this construction requires the existence of an inner product.

Notice that this is indeed a bounded linear map, where boundedness comes from the Cauchy-Schwarz inequality

$$|\langle \mathbf{w}, \mathbf{v} \rangle| \leq \|\mathbf{v}\|_H \|\mathbf{w}\|_H.$$

Therefore, ϕ taking $\mathbf{v} \mapsto \phi_{\mathbf{v}}$ is a map $\phi : H \rightarrow H^*$.

Using this simple construction, we have managed to produce a lot of members of the dual. Are there any more things in the dual? The answer is no, and this is given by the Riesz representation theorem.

Proposition (Riesz representation theorem). Let H be a Hilbert space. Then $\phi : H \rightarrow H^*$ defined by $\mathbf{v} \mapsto \langle \cdot, \mathbf{v} \rangle$ is an isometric anti-isomorphism, i.e. it is isometric, bijective and

$$\phi(\lambda \mathbf{v} + \mu \mathbf{w}) = \bar{\lambda} \phi(\mathbf{v}) + \bar{\mu} \phi(\mathbf{w}).$$

Proof. We first prove all the easy bits, namely everything but surjectivity.

- To show injectivity, if $\phi_{\mathbf{v}} = \phi_{\mathbf{u}}$, then $\langle \mathbf{w}, \mathbf{v} \rangle = \langle \mathbf{w}, \mathbf{u} \rangle$ for all \mathbf{w} by definition. So $\langle \mathbf{w}, \mathbf{v} - \mathbf{u} \rangle = 0$ for all \mathbf{w} . In particular, $\langle \mathbf{v} - \mathbf{w}, \mathbf{v} - \mathbf{w} \rangle = 0$. So $\mathbf{v} - \mathbf{w} = 0$.
- To show that it is an anti-homomorphism, let $\mathbf{v}, \mathbf{w}, \mathbf{y} \in H$ and $\lambda, \mu \in \mathbb{F}$. Then

$$\phi_{\lambda \mathbf{v} + \mu \mathbf{w}}(\mathbf{y}) = \langle \mathbf{y}, \lambda \mathbf{v} + \mu \mathbf{w} \rangle = \bar{\lambda} \langle \mathbf{y}, \mathbf{v} \rangle + \bar{\mu} \langle \mathbf{y}, \mathbf{w} \rangle = \bar{\lambda} \phi_{\mathbf{v}}(\mathbf{y}) + \bar{\mu} \phi_{\mathbf{w}}(\mathbf{y}).$$

- To show it is isometric, let $\mathbf{v}, \mathbf{w} \in H$ and $\|\mathbf{w}\|_H = 1$. Then

$$|\phi_{\mathbf{v}}(\mathbf{w})| = |\langle \mathbf{w}, \mathbf{v} \rangle| \leq \|\mathbf{w}\|_H \|\mathbf{v}\|_H = \|\mathbf{v}\|_H.$$

Hence, for all \mathbf{v} , $\|\phi_{\mathbf{v}}\|_{H^*} \leq \|\mathbf{v}\|_H$ for all $\mathbf{v} \in H$. To show $\|\phi_{\mathbf{v}}\|_{H^*}$ is exactly $\|\mathbf{v}\|_H$, it suffices to note that

$$|\phi_{\mathbf{v}}(\mathbf{v})| = \langle \mathbf{v}, \mathbf{v} \rangle = \|\mathbf{v}\|_H^2.$$

So $\|\phi_{\mathbf{v}}\|_{H^*} \geq \|\mathbf{v}\|_H^2 / \|\mathbf{v}\|_H = \|\mathbf{v}\|_H$.

Finally, we show surjectivity. Let $\xi \in H^*$. If $\xi = 0$, then $\xi = \phi_{\mathbf{0}}$.

Otherwise, suppose $\xi \neq 0$. The idea is that $(\ker \xi)^\perp$ is one-dimensional, and then the \mathbf{v} we are looking for will be an element in this complement. So we arbitrarily pick one, and then scale it appropriately.

We now write out the argument carefully. First, we note that since ξ is continuous, $\ker \xi$ is closed, since it is the inverse image of the closed set $\{0\}$. So $\ker \xi$ is complete, and thus we have

$$H = \ker \xi \oplus (\ker \xi)^\perp.$$

The next claim is that $\dim(\ker \xi) = 1$. This is an immediate consequence of the first isomorphism theorem, whose proof is the usual one, but since we didn't prove that, we will run the argument manually.

We pick any two elements $\mathbf{v}_1, \mathbf{v}_2 \in (\ker \xi)^\perp$. Then we can always find some λ, μ not both zero such that

$$\lambda \xi(\mathbf{v}_1) + \mu \xi(\mathbf{v}_2) = 0.$$

So $\lambda \mathbf{v}_1 + \mu \mathbf{v}_2 \in \ker \xi$. But they are also in $(\ker \xi)^\perp$ by linearity. Since $\ker \xi$ and $(\ker \xi)^\perp$ have trivial intersection, we deduce that $\lambda \mathbf{v}_1 + \mu \mathbf{v}_2 = 0$. Thus, any two vectors in $(\ker \xi)^\perp$ are dependent. Since $\xi \neq 0$, we know that $\ker \xi$ has dimension 1.

Now pick any $\mathbf{v} \in (\ker \xi)^\perp$ such that $\xi(\mathbf{v}) \neq 0$. By scaling it appropriately, we can obtain a \mathbf{v} such that

$$\xi(\mathbf{v}) = \langle \mathbf{v}, \mathbf{v} \rangle.$$

Finally, we show that $\xi = \phi_{\mathbf{v}}$. To prove this, let $\mathbf{w} \in H$. We decompose \mathbf{w} using the previous theorem to get

$$\mathbf{w} = \alpha \mathbf{v} + \mathbf{w}_0$$

for some $\mathbf{w}_0 \in \ker \xi$ and $\alpha \in \mathbb{F}$. Note that by definition of $(\ker \xi)^\perp$, we know that $\langle \mathbf{w}_0, \mathbf{v} \rangle = 0$. Hence we know that

$$\begin{aligned} \xi(\mathbf{w}) &= \xi(\alpha \mathbf{v} + \mathbf{w}_0) = \xi(\alpha \mathbf{v}) = \alpha \xi(\mathbf{v}) \\ &= \alpha \langle \mathbf{v}, \mathbf{v} \rangle = \langle \alpha \mathbf{v}, \mathbf{v} \rangle = \langle \alpha \mathbf{v} + \mathbf{w}_0, \mathbf{v} \rangle = \langle \mathbf{w}, \mathbf{v} \rangle. \end{aligned}$$

Since \mathbf{w} was arbitrary, we are done. \square

Using this proposition twice, we know that all Hilbert spaces are reflexive, i.e. $H \cong H^{**}$.

We now return to the proof of the proposition we claimed at the beginning.

Proposition. For $f \in C(S^1)$, defined, for each $k \in \mathbb{Z}$,

$$\hat{f}(k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ikx} f(x) \, dx.$$

The partial sums are then defined as

$$S_N(f)(x) = \sum_{n=-N}^N e^{inx} \hat{f}(k).$$

Then we have

$$\lim_{N \rightarrow \infty} \frac{1}{2\pi} \int_{-\pi}^{\pi} |f(x) - S_N(f)(x)|^2 \, dx = 0.$$

Proof. Consider the following Hilbert space $L^2(S^1)$ defined as the completion of $C_{\mathbb{C}}(S^1)$ under the inner product

$$\langle f, g \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \bar{g}(x) \, dx,$$

Consider the closed subspace

$$U_N = \text{span}\{e^{inx} : |n| \leq N\}.$$

Then in fact S_N defined above by

$$S_N(f)(x) = \sum_{n=-N}^N e^{-inx} \hat{f}(k)$$

is the projection operator onto U_N . This is since we have the orthonormal condition

$$\langle e^{inx}, e^{-imx} \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{inx} e^{-imx} dx = \begin{cases} 1 & n = m \\ 0 & n \neq m \end{cases}$$

Hence it is easy to check that if $f \in U_N$, say $f = \sum_{n=-N}^N a_n e^{inx}$, then $S_N f = f$ since

$$S_N(f) = \sum_{n=-N}^N \hat{f}(k) e^{-inx} = \sum_{n=-N}^N \langle f, e^{inx} \rangle e^{-inx} = \sum_{n=-N}^N a_n e^{-inx} = f$$

using the orthogonality relation. But if $f \in U_N^\perp$, then

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-inx} f(x) dx = 0$$

for all $|n| < N$. So $S_N(f) = 0$. So this is indeed a projection map.

In particular, we will use the fact that projection maps have norms ≤ 1 . Hence for any $P(x)$, we have

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} |S_N(f)(x) - S_N(P)(x)|^2 dx \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} |f(x) - P(x)|^2 dx$$

Now consider the algebra \mathcal{A} generated $\{e^{inx} : n \in \mathbb{Z}\}$. Notice that \mathcal{A} separates points and is closed under complex conjugation. Also, for every $x \in S^1$, there exists $f \in \mathcal{A}$ such that $f(x) \neq 0$ (using, say $f(x) = e^{ix}$). Hence, by Stone-Weierstrass theorem, $\bar{\mathcal{A}} = C_{\mathbb{C}}(S^1)$, i.e. for every $f \in C_{\mathbb{C}}(S^1)$ and $\varepsilon > 0$, there exists a polynomial P of e^{ix} and e^{-ix} such that

$$\|P - f\| < \varepsilon.$$

We are almost done. We now let $N > \deg P$ be a large number. Then in particular, we have $S_N(P) = P$. Then

$$\begin{aligned} \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |S_N(f) - f|^2 dx \right)^{\frac{1}{2}} &\leq \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |S_N(f) - S_N(P)|^2 dx \right)^{\frac{1}{2}} \\ &\quad + \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |S_N(P) - P|^2 dx \right)^{\frac{1}{2}} \\ &\quad + \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |P - f|^2 dx \right)^{\frac{1}{2}} \\ &\leq \varepsilon + 0 + \varepsilon \\ &= 2\varepsilon. \end{aligned}$$

So done. □

4.3 Orthonormal systems and basis

Definition (Orthonormal system). Let E be a Euclidean space. A set of unit vectors $\{\mathbf{e}_\alpha\}_{\alpha \in A}$ is called an *orthonormal system* if $\langle \mathbf{e}_\alpha, \mathbf{e}_\beta \rangle = 0$ if $\alpha \neq \beta$.

We want to define a “basis” in an infinite dimensional vector space. The idea is that these should be orthonormal systems “big enough” to span everything. In finite-dimensions, this was easy, since there is the notion of dimension — if we have n dimensions, then we just take an orthonormal system of n vectors, and we are done.

If we have infinite dimensions, this is trickier. If we have many many dimensions and vectors, we can keep adding things to our orthonormal system, but we might never get to such a “basis”, if our “basis” has to be uncountable. Hence we have the idea of “maximality”.

Definition (Maximal orthonormal system). Let E be a Euclidean space. An orthonormal space is called *maximal* if it cannot be extended to a strictly larger orthonormal system.

By Zorn’s lemma, a maximal orthonormal system always exists. We will later see that in certain nice cases, we can construct a maximal orthonormal system directly, without appealing to Zorn’s lemma. The advantage of an explicit construction is that we will understand our system much more.

One important thing we would like to do is given an orthonormal system, decide whether it is maximal. In general, this is difficult, and Zorn’s lemma is completely useless.

Now suppose we are nicer and have a Hilbert space. What we would like to say is that if we have a maximal orthonormal system, then its span is the whole space H . However, this doesn’t really work. The span of a set S only allows us to take finite linear combinations, but by completeness of H , we want to have the infinite sums, i.e. the limits as well. So what we really have is the following.

Proposition. Let H be a Hilbert space. Let S be a maximal orthonormal system. Then $\overline{\text{span } S} = H$.

While this might seem difficult to prove at first, it turns out the proof is pretty short and simple.

Proof. Recall that $S^\perp = (\overline{\text{span } S})^\perp$. Since H is a Hilbert space, we have

$$H = \overline{\text{span } S} \oplus (\overline{\text{span } S})^\perp = \overline{\text{span } S} \oplus S^\perp.$$

Since S is maximal, $S^\perp = \{0\}$. So done. □

How about the converse? It is also true. In fact, it is true even for Euclidean spaces, and the proof is easy.

Proposition. Let E be Euclidean, and let S be an orthonormal system. If $\overline{\text{span } S} = E$, then S is maximal.

Proof.

$$S^\perp = (\overline{\text{span } S})^\perp = E^\perp = \{0\}. \quad \square$$

So in a Hilbert space, we have an if and only if condition — a system is maximal if and only if the closure of the span is everything. In other words, given any vector $\mathbf{v} \in H$, we can find a sequence \mathbf{v}_i in the span of the maximal system that converges to \mathbf{v} . This sequence is clearly not unique, since we can just add a random term to the first item.

However, we can do something better. Consider our space ℓ_2 , and the element $(1, \frac{1}{2}, \frac{1}{4}, \dots)$. There is a very natural way to write this as the limit of the sequence:

$$(1, 0, 0, \dots), \left(1, \frac{1}{2}, 0, \dots\right), \left(1, \frac{1}{2}, \frac{1}{4}, 0, \dots\right), \dots$$

What we are doing is that we are truncating the element at the n th component for each n . Alternatively, the n th term is what we get when we project our \mathbf{v} onto the space spanned by the first n “basis” vectors. This is a nice and natural way to produce the sequence.

Definition (Hilbert space basis). Let H be a Hilbert space. A maximal orthonormal system is called a *Hilbert space basis*.

Recall that at the beginning, we said we needed Zorn’s lemma to get an orthonormal system. In many cases, we can find a basis *without* using Zorn’s lemma. This relies on the Gram-Schmidt procedure.

Proposition. Let $\{\mathbf{x}_i\}_{i=1}^n$, $n \in \mathbb{N}$ be linearly independent. Then there exists $\{\mathbf{e}_i\}_{i=1}^n$ such that $\{\mathbf{e}_i\}_{i=1}^n$ is an orthonormal system and

$$\text{span}\{\mathbf{x}_1, \dots, \mathbf{x}_j\} = \text{span}\{\mathbf{e}_1, \dots, \mathbf{e}_j\}$$

for all $j \leq n$.

Proof. Define \mathbf{e}_1 by

$$\mathbf{e}_1 = \frac{\mathbf{x}_1}{\|\mathbf{x}_1\|}.$$

Assume we have defined $\{\mathbf{e}_i\}_{i=1}^j$ orthonormal such that

$$\text{span}\{\mathbf{x}_1, \dots, \mathbf{x}_j\} = \text{span}\{\mathbf{e}_1, \dots, \mathbf{e}_j\}.$$

Then by linear independence, we know that

$$\mathbf{x}_{j+1} \notin \text{span}\{\mathbf{x}_1, \dots, \mathbf{x}_j\} = \text{span}\{\mathbf{e}_1, \dots, \mathbf{e}_j\} = F_j.$$

We now define

$$\tilde{\mathbf{x}}_{j+1} = \mathbf{x}_{j+1} - P_{F_j}(\mathbf{x}_{j+1}),$$

where P_{F_j} is the projection onto F_j given by

$$P_{F_j} = \sum_{i=1}^j \langle \mathbf{x}, \mathbf{e}_i \rangle \mathbf{e}_i.$$

Since F_j is a closed, finite subspace, we know that

$$\mathbf{x}_{j+1} - P_{F_j} \mathbf{x}_{j+1} \perp F_j.$$

Thus

$$\mathbf{e}_{j+1} = \frac{\tilde{\mathbf{x}}_{j+1}}{\|\tilde{\mathbf{x}}_{j+1}\|}$$

is the right choice. We can also write this in full as

$$\mathbf{e}_{j+1} = \frac{\mathbf{x}_{j+1} - \sum_{i=1}^j \langle \mathbf{x}_j \mathbf{e}_j \rangle \mathbf{e}_i}{\|\mathbf{x}_{j+1} - \sum_{i=1}^j \langle \mathbf{x}_j \mathbf{e}_j \rangle \mathbf{e}_i\|}.$$

So done. \square

Note that projection into the first n basis is exactly what we did when we wrote an element in ℓ_2 as the limit of the sequence.

This is a helpful result, since it is a constructive way of producing orthonormal systems. So if we are handed a set of vectors, we can just apply this result, plug our vectors into this ugly formula, and get a result. Of course, we want to apply this to infinite spaces.

Proposition. Let H be separable, i.e. there is an infinite set $\{\mathbf{y}_i\}_{i \in \mathbb{N}}$ such that

$$\overline{\text{span}\{\mathbf{y}_i\}} = H.$$

Then there exists a countable basis for $\text{span}\{\mathbf{y}_i\}$.

Proof. We find a subset $\{\mathbf{y}_{i_j}\}$ such that $\text{span}\{\mathbf{y}_i\} = \text{span}\{\mathbf{y}_{i_j}\}$ and $\{\mathbf{y}_{i_j}\}$ are independent. This is easy to do since we can just throw away the useless dependent stuff. At this point, we do Gram-Schmidt, and done. \square

Example. Consider $H = \ell_2$ and the sequence $\{\mathbf{e}_i\}_{i \in \mathbb{N}}$ be defined by

$$\mathbf{e}_i = (0, 0, \dots, 0, 1, 0, \dots),$$

with the zero in the i th column.

Note that $\mathbf{x} \perp \{\mathbf{e}_i\}_{i \in \mathbb{N}}$ if and only if each component is zero, i.e. $\mathbf{x} = \mathbf{0}$. So $\{\mathbf{e}_i\}$ is maximal, and hence a basis.

Example. Consider $H = L^2$, the completion of $C(S^1)$ under the L^2 norm, i.e.

$$\langle f, g \rangle = \int_{-\pi}^{\pi} f \bar{g} \, dx.$$

Trigonometric polynomials are dense in $C(S^1)$ with respect to the supremum norm due to Stone-Weierstrass. So in fact $\text{span}\left\{\frac{1}{\sqrt{2\pi}}e^{inx}\right\}$ for $n \in \mathbb{N}$ is dense in $C(S^1)$. Hence it is dense in $C(S^1)$ under the L^2 norm since convergence under the supremum norm implies convergence under L^2 . In particular, it is dense in the L^2 space since L^2 is the completion of $C(S^1)$. Moreover, this set is orthonormal in $C(S^1)$ under the L^2 norm. So $\left\{\frac{1}{\sqrt{2\pi}}e^{inx}\right\}$ is a basis for L^2 .

Note that in these two examples, we have exhibited two different ways of constructing a basis. In the first case, we showed that it is maximal directly. In the second case, we show that its span is a dense subset of the space. By our proposition, these are equivalent and valid ways of proving that it is a basis.

4.4 The isomorphism with ℓ_2

We ended the previous section with two examples. Both of them are Hilbert spaces, and both have countable basis. Is there any way we can identify the both? This is a reasonable thing to ask. If we are given a Hilbert space H of finite dimension $\dim H = n$, then we know that H is indeed isomorphic to \mathbb{R}^n (or \mathbb{C}^n) with the Euclidean norm. In some sense ℓ_2 is just an “infinite version” of \mathbb{R}^n . So we might expect all other Hilbert spaces with countable dimension to be isomorphic to ℓ_2 .

Recall that if we have a finite-dimensional Hilbert space H with $\dim H = n$, and an orthonormal basis $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$, then each $\mathbf{x} \in H$ can be written as

$$\mathbf{x} = \sum_{i=1}^n \langle \mathbf{x}, \mathbf{e}_i \rangle \mathbf{e}_i,$$

and

$$\|\mathbf{x}\|^2 = \sum_{i=1}^n |\langle \mathbf{x}, \mathbf{e}_i \rangle|^2.$$

Thus H is isomorphic to ℓ_2^n , the space \mathbb{R}^n with the Euclidean norm, via the map

$$\mathbf{x} \mapsto (\langle \mathbf{x}, \mathbf{e}_1 \rangle, \dots, \langle \mathbf{x}, \mathbf{e}_n \rangle).$$

Can we push this to the infinite dimensional case? Yes. We will have to replace our finite sum $\sum_{i=1}^n$ with an infinite sum. Of course, with an infinite sum, we need to make sure things converge. This is guaranteed by Bessel’s inequality.

Lemma (Bessel’s inequality). Let E be Euclidean and $\{\mathbf{e}_i\}_{i=1}^N$ with $N \in \mathbb{N} \cup \{\infty\}$ an orthonormal system. For any $\mathbf{x} \in E$, define $x_i = \langle \mathbf{x}, \mathbf{e}_i \rangle$. Then for any $j \leq N$, we have

$$\sum_{i=1}^j |x_i|^2 \leq \|\mathbf{x}\|^2.$$

Proof. Consider the case where j is finite first. Define

$$F_j = \text{span}\{\mathbf{e}_1, \dots, \mathbf{e}_j\}.$$

This is a finite dimensional subspace of E . Hence an orthogonal projection P_{F_j} exists. Moreover, we have an explicit formula for this:

$$P_{F_j} = \sum_{i=1}^j \langle \mathbf{x}, \mathbf{e}_i \rangle \mathbf{e}_i.$$

Thus

$$\sum_{i=1}^j |x_i|^2 = \|P_{F_j} \mathbf{x}\|^2 \leq \|\mathbf{x}\|^2$$

since we know that $\|P_{F_j}\| \leq 1$. Taking the limit as $j \rightarrow \infty$ proves the case for infinite j . \square

The only thing we required in the proof is for the space to be Euclidean. This is since we are talking about the sum

$$\sum_{i=1}^{\infty} |x_i|^2,$$

and this is a sum of numbers. However, if we want to investigate the sum

$$\mathbf{x} = \sum_{i=1}^{\infty} \langle \mathbf{x}, \mathbf{e}_i \rangle \mathbf{e}_i,$$

then we'd better require the space to be Hilbert, so that the sum has something to converge to.

Proposition. Let H be a separable Hilbert space, with a countable basis $\{\mathbf{e}_i\}_{i=1}^N$, where $N \in \mathbb{N} \cup \{\infty\}$. Let $\mathbf{x}, \mathbf{y} \in H$ and

$$x_i = \langle \mathbf{x}, \mathbf{e}_i \rangle, \quad y_i = \langle \mathbf{y}, \mathbf{e}_i \rangle.$$

Then

$$\mathbf{x} = \sum_{i=1}^N x_i \mathbf{e}_i, \quad \mathbf{y} = \sum_{i=1}^N y_i \mathbf{e}_i,$$

and

$$\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{i=1}^N x_i \bar{y}_i.$$

Moreover, the sum converges absolutely.

Proof. We only need to consider the case $N = \infty$. Otherwise, it is just finite-dimensional linear algebra.

First, note that our expression is written as an infinite sum. So we need to make sure it converges. We define the partial sums to be

$$\mathbf{s}_n = \sum_{i=1}^n x_i \mathbf{e}_i.$$

We want to show $\mathbf{s}_n \rightarrow \mathbf{x}$. By Bessel's inequality, we know that

$$\sum_{i=1}^{\infty} |x_i|^2 \leq \|\mathbf{x}\|^2.$$

In particular, the sum is bounded, and hence converges.

For any $m < n$, we have

$$\|\mathbf{s}_n - \mathbf{s}_m\| = \sum_{i=m+1}^n |x_i|^2 \leq \sum_{i=m+1}^{\infty} |x_i|^2.$$

As $m \rightarrow \infty$, the series must go to 0. Thus $\{\mathbf{s}_n\}$ is Cauchy. Since H is Hilbert, \mathbf{s}_n converges, say

$$\mathbf{s}_n \rightarrow \mathbf{s} = \sum_{i=1}^{\infty} x_i \mathbf{e}_i.$$

Now we want to prove that this sum is indeed \mathbf{x} itself. Note that so far in the proof, we have *not* used the fact that $\{\mathbf{e}_i\}$ is a basis. We just used the fact that it is orthogonal. Hence we should use this now. We notice that

$$\langle \mathbf{s}, \mathbf{e}_i \rangle = \lim_{n \rightarrow \infty} \langle \mathbf{s}_n, \mathbf{e}_i \rangle = \lim_{n \rightarrow \infty} \sum_{j=1}^n x_j \langle \mathbf{e}_j, \mathbf{e}_i \rangle = x_i.$$

Hence we know that

$$\langle \mathbf{x} - \mathbf{s}, \mathbf{e}_i \rangle = 0.$$

for all i . So $\mathbf{x} - \mathbf{s}$ is perpendicular to all \mathbf{e}_i . Since $\{\mathbf{e}_i\}$ is a basis, we must have $\mathbf{x} - \mathbf{s} = 0$, i.e. $\mathbf{x} = \mathbf{s}$.

To show our formula for the inner product, we can compute

$$\begin{aligned} \langle \mathbf{x}, \mathbf{y} \rangle &= \lim_{n \rightarrow \infty} \left\langle \sum_{i=1}^n x_i \mathbf{e}_i, \sum_{j=1}^n y_j \mathbf{e}_j \right\rangle \\ &= \lim_{n \rightarrow \infty} \sum_{i,j=1}^n x_i \bar{y}_j \langle \mathbf{e}_i, \mathbf{e}_j \rangle \\ &= \lim_{n \rightarrow \infty} \sum_{i,j=1}^n x_i \bar{y}_j \delta_{ij} \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n x_i \bar{y}_i \\ &= \sum_{i=1}^{\infty} x_i \bar{y}_i. \end{aligned}$$

Note that we *know* the limit exists, since the continuity of the inner product ensures the first line is always valid.

Finally, to show absolute convergence, note that for all finite j , we have

$$\sum_{i=1}^j |x_i \bar{y}_i| \leq \sqrt{\sum_{i=1}^n |x_i|^2} \sqrt{\sum_{i=1}^n |y_i|^2} \leq \|\mathbf{x}\| \|\mathbf{y}\|.$$

Since this is a uniform bound for any j , the sum converges absolutely. \square

Note that in the case of $\mathbf{x} = \mathbf{y}$, our formula for the inner product gives

$$\|\mathbf{x}\|^2 = \sum_{i=1}^N |x_i|^2.$$

This is known as *Parseval's equality*

What this proposition gives us is that given any separable Hilbert space, we can find “coordinates” for it, and in terms of these coordinates, our inner product and hence norm all act like ℓ_2 . In particular, we have the map

$$\mathbf{x} \mapsto \{\langle \mathbf{x}, \mathbf{e}_i \rangle\}_{i=1}^N$$

that takes H into ℓ_2 . This is injective since by Parseval's equality, if the image of \mathbf{x} is 0, then $\|\mathbf{x}\|^2 = \sum 0 = 0$. So $\mathbf{x} = 0$.

This is good, but not good enough. We want the map to be an isomorphism. Hence, we need to show it is surjective. In other words, *every* element in ℓ_2 is obtained. This is a theorem by Riesz and Fisher, and is in fact easy to prove, since there is an obvious candidate for the preimage of any $\{x_i\}_{i \in \mathbb{N}}$.

Proposition. Let H be a separable Hilbert space with orthonormal basis $\{\mathbf{e}_i\}_{i \in \mathbb{N}}$. Let $\{a_i\}_{i \in \mathbb{N}} \in \ell_2(\mathbb{C})$. Then there exists an $\mathbf{x} \in H$ with $\langle \mathbf{x}, \mathbf{e}_i \rangle = a_i$. Moreover, this \mathbf{x} is exactly

$$\mathbf{x} = \sum_{i=1}^{\infty} x_i \mathbf{e}_i.$$

Proof. The only thing we need to show is that this sum converges. For any $n \in \mathbb{N}$, define

$$\mathbf{s}_n = \sum_{i=1}^n a_i \mathbf{e}_i \in H.$$

For $m < n$, we have

$$\|\mathbf{s}_n - \mathbf{s}_m\|^2 = \sum_{m+1}^n |a_i|^2 \rightarrow 0$$

as $m \rightarrow \infty$ because $\{a_i\} \in \ell^2$. Hence \mathbf{s}_n is Cauchy and as such converges to \mathbf{x} . Obviously, we have

$$\langle \mathbf{x}, \mathbf{e}_i \rangle = \lim_{n \rightarrow \infty} \sum_{j=1}^n a_j \langle \mathbf{e}_j, \mathbf{e}_i \rangle = a_i.$$

So done. □

This means we have a isomorphism between ℓ_2 and H . Moreover, this is continuous and in fact isometric. So this is a very strong result. This says all separable Hilbert spaces *are* ℓ_2 .

4.5 Operators

We are going to look at operators on Hilbert spaces. For example, we would like to see how differential operators behave on spaces of differentiable functions.

In this section, we will at least require the space to be Banach. So let X be a Banach space over \mathbb{C} . We will consider $\mathcal{B}(X) = \mathcal{B}(X, X)$, the vector space of bounded linear maps from X to itself. We have seen in the example sheets that $\mathcal{B}(X)$ is a unital Banach algebra, i.e. it forms a complete algebra with composition as multiplication. Our goal is to generalize some considerations in finite dimensions such as eigenvectors and eigenvalues.

Definition (Spectrum and resolvent set). Let X be a Banach space and $T \in \mathcal{B}(X)$, we define the *spectrum* of T , denoted by $\sigma(T)$ by

$$\sigma(t) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not invertible}\}.$$

The *resolvent set*, denoted by $\rho(T)$, is

$$\rho(t) = \mathbb{C} \setminus \sigma(T).$$

Note that if $T - \lambda$ is bijective, then by the inverse mapping theorem, we know it has a bounded inverse. So if $\lambda \in \sigma(T)$, then either $T - \lambda I$ is not injective, or it is not surjective. In other words, $\ker(T - \lambda I) \neq \{0\}$ or $\text{im}(T - \lambda I) \neq X$. In finite dimensions, these are equivalent by, say, the rank-nullity theorem, but in general, they are not.

Example. Consider the shift operator $s : \ell_\infty \rightarrow \ell_\infty$ defined by

$$(a_1, a_2, a_3, \dots) \mapsto (0, a_1, a_2, \dots).$$

Then this is injective but not surjective.

Now if $\lambda \in \rho(T)$, i.e. $T - \lambda I$ is invertible, then $(T - \lambda I)^{-1}$ is automatically bounded by the inverse mapping theorem. This is why we want to work with Banach spaces.

Definition (Resolvent). Let X be a Banach space. The *resolvent* is the map $R : \rho(T) \rightarrow \mathcal{B}(X)$ given by

$$\lambda \mapsto (T - \lambda I)^{-1}.$$

Definition (Eigenvalue). We say λ is an *eigenvalue* of T if $\ker(T - \lambda I) \neq \{0\}$.

Definition (Point spectrum). Let X be a Banach space. The *point spectrum* is

$$\sigma_p(T) = \{\lambda \in \mathbb{C} : \lambda \text{ is an eigenvalue of } T\}.$$

Obviously, $\sigma_p(T) \subseteq \sigma(T)$, but they are in general not equal.

Definition (Approximate point spectrum). Let X be a Banach space. The *approximate point spectrum* is defined as

$$\sigma_{ap}(X) = \{\lambda \in \mathbb{C} : \exists \{\mathbf{x}_n\} \subseteq X : \|\mathbf{x}_n\|_X = 1 \text{ and } \|(T - \lambda I)\mathbf{x}_n\|_X \rightarrow 0\}.$$

Again, we have

$$\sigma_p(T) \subseteq \sigma_{ap}(T) \subseteq \sigma(T).$$

The last inclusion follows from the fact that if an inverse exists, then the inverse is bounded.

An important characterization of the spectrum is the following theorem:

Theorem. Let X be a Banach space, $T \in \mathcal{B}(X)$. Then $\sigma(T)$ is a non-empty, closed subset of

$$\{\lambda \in \mathbb{C} : |\lambda| \leq \|T\|_{\mathcal{B}(X)}\}.$$

In finite dimensions, this in particular implies the existence of eigenvalues, since the spectrum is equal to the point spectrum. Notice this is only true for vector spaces over \mathbb{C} , as we know from linear algebra.

To prove this theorem, we will first prove two lemmas.

Lemma. Let X be a Banach space, $T \in \mathcal{B}(X)$ and $\|T\|_{\mathcal{B}(X)} < 1$. Then $I - T$ is invertible.

Proof. To prove it is invertible, we construct an explicit inverse. We want to show

$$(I - T)^{-1} = \sum_{i=0}^{\infty} T^i.$$

First, we check the right hand side is absolutely convergent. This is since

$$\sum_{i=0}^{\infty} \|T^i\|_{\mathcal{B}(X)} \leq \sum_{i=0}^{\infty} \|T\|_{\mathcal{B}(X)}^i \leq \frac{1}{1 - \|T\|_{\mathcal{B}(X)}} < \infty.$$

Since X is Banach, and hence $\mathcal{B}(X)$ is Banach, the limit is well-defined. Now it is easy to check that

$$\begin{aligned} (I - T) \sum_{i=0}^{\infty} T^i &= (I - T)(I + T + T^2 + \cdots) \\ &= I + (T - T) + (T^2 - T^2) + \cdots \\ &= I. \end{aligned}$$

Similarly, we have

$$\left(\sum_{i=1}^{\infty} T^i \right) (I - T) = I. \quad \square$$

Lemma. Let X be a Banach space, $S_1 \in \mathcal{B}(X)$ be invertible. Then for all $S_2 \in \mathcal{B}(X)$ such that

$$\|S_1^{-1}\|_{\mathcal{B}(X)} \|S_1 - S_2\|_{\mathcal{B}(X)} < 1,$$

S_2 is invertible.

This is some sort of an “openness” statement for the invertible bounded linear maps, since if S_1 is invertible, then any “nearby” bounded linear map is also invertible.

Proof. We can write

$$S_2 = S_1(I - S_1^{-1}(S_1 - S_2)).$$

Since

$$\|S_1^{-1}(S_1 - S_2)\|_{\mathcal{B}(X)} \leq \|S_1^{-1}\|_{\mathcal{B}(X)} \|S_1 - S_2\|_{\mathcal{B}(X)} < 1$$

by assumption, by the previous lemma, $(I - S_1^{-1}(S_1 - S_2))^{-1}$ exists. Therefore the inverse of S_2 is

$$S_2^{-1} = (I - S_1^{-1}(S_1 - S_2))^{-1} S_1^{-1}. \quad \square$$

We can now return to prove our original theorem.

Theorem. Let X be a Banach space, $T \in \mathcal{B}(X)$. Then $\sigma(T)$ is a non-empty, closed subset of

$$\{\lambda \in \mathbb{C} : |\lambda| \leq \|T\|_{\mathcal{B}(X)}\}.$$

Note that it is not hard to prove that it is closed and a subset of $\{\lambda \in \mathbb{C} : |\lambda| \leq \|T\|_{\mathcal{B}(X)}\}$. The hard part is to prove it is non-empty.

Proof. We first prove the closedness of the spectrum. It suffices to prove that the resolvent set $\rho(T) = \mathbb{C} \setminus \sigma(T)$ is open, by the definition of closedness.

Let $\lambda \in \rho(T)$. By definition, $S_1 = T - \lambda I$ is invertible. Define $S_2 = T - \mu I$. Then

$$\|S_1 - S_2\|_{\mathcal{B}(X)} = \|(T - \lambda I) - (T - \mu I)\|_{\mathcal{B}(X)} = |\lambda - \mu|.$$

Hence if $|\lambda - \mu|$ is sufficiently small, then $T - \mu I$ is invertible by the above lemma. Hence $\mu \in \rho(T)$. So $\rho(T)$ is open.

To show $\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| \leq \|T\|_{\mathcal{B}(X)}\}$ is equivalent to showing

$$\{\lambda \in \mathbb{C} : |\lambda| > \|T\|_{\mathcal{B}(X)}\} \subseteq \mathbb{C} \setminus \sigma(T) = \rho(T).$$

Suppose $|\lambda| > \|T\|$. Then $I - \lambda^{-1}T$ is invertible since

$$\|\lambda^{-1}T\|_{\mathcal{B}(X)} = \lambda^{-1}\|T\|_{\mathcal{B}(X)} < 1.$$

Therefore, $(I - \lambda^{-1}T)^{-1}$ exists, and hence

$$(\lambda I - T)^{-1} = \lambda^{-1}(I - \lambda^{-1}T)^{-1}$$

is well-defined. Therefore $\lambda I - T$, and hence $T - \lambda I$ is invertible. So $\lambda \in \rho(T)$.

Finally, we need to show it is non-empty. How did we prove it in the case of finite-dimensional vector spaces? In that case, it ultimately boiled down to the fundamental theorem of algebra. And how did we prove the fundamental theorem of algebra? We said that if $p(x)$ is a polynomial with no roots, then $\frac{1}{p(x)}$ is bounded and entire, hence constant.

We are going to do the same proof. We look at $\frac{1}{T - \lambda I}$ as a function of λ . If $\sigma(T) = \emptyset$, then this is an everywhere well-defined function. We show that this is entire and bounded, and hence by “Liouville’s theorem”, it must be constant, which is impossible (in the finite-dimensional case, we would have inserted a det there).

So suppose $\sigma(T) = \emptyset$, and consider the function $R : \mathbb{C} \rightarrow \mathcal{B}(X)$, given by $R(\lambda) = (T - \lambda I)^{-1}$.

We first show this is entire. This, by definition, means R is given by a power series near any point $\lambda_0 \in \mathbb{C}$. Fix such a point. Then as before, we can expand

$$\begin{aligned} T - \lambda I &= (T - \lambda_0 I) \left[I - (T - \lambda_0 I)^{-1} \left((T - \lambda_0 I) - (T - \lambda I) \right) \right] \\ &= (T - \lambda_0 I) \left[I - (\lambda - \lambda_0)(T - \lambda_0 I)^{-1} \right]. \end{aligned}$$

Then for $(\lambda - \lambda_0)$ small, we have

$$\begin{aligned} (T - \lambda I)^{-1} &= \left(\sum_{i=0}^{\infty} (\lambda - \lambda_0)^i (T - \lambda_0 I)^{-i} \right) (T - \lambda_0 I)^{-1} \\ &= \sum_{i=0}^{\infty} (\lambda - \lambda_0)^i (T - \lambda_0 I)^{-i-1}. \end{aligned}$$

So this is indeed given by an absolutely convergent power series near λ_0 .

Next, we show R is bounded, i.e.

$$\sup_{\lambda \in \mathbb{C}} \|R(\lambda)\|_{\mathcal{B}(X)} < \infty.$$

It suffices to prove this for λ large. Note that we have

$$(T - \lambda I)^{-1} = \lambda^{-1}(\lambda^{-1}T - I)^{-1} = -\lambda^{-1} \sum_{i=0}^{\infty} \lambda^{-i} T^i.$$

Hence we get

$$\begin{aligned} \|(\lambda I - T)^{-1}\|_{\mathcal{B}(X)} &\leq |\lambda|^{-1} \sum_{i=0}^{\infty} |\lambda|^{-i} \|T^i\|_{\mathcal{B}(X)} \\ &\leq |\lambda|^{-1} \sum_{i=0}^{\infty} (|\lambda|^{-1} \|T\|_{\mathcal{B}(X)})^i \\ &\leq \frac{1}{|\lambda| - \|T\|_{\mathcal{B}(X)}}, \end{aligned}$$

which tends to 0 as $|\lambda| \rightarrow \infty$. So it is bounded.

By “Liouville’s theorem”, $R(\lambda)$ is constant, which is clearly a contradiction since $R(\lambda) \neq R(\mu)$ for $\lambda \neq \mu$. \square

Of course, to do this properly, we need a version of Liouville’s theorem for Banach-space valued functions as opposed to complex-valued functions. So let’s prove this.

Proposition (Liouville’s theorem for Banach space-valued analytic function). Let X be a Banach space, and $F : \mathbb{C} \rightarrow X$ be entire (in the sense that F is given by an absolutely convergent power series in some neighbourhood of any point) and norm bounded, i.e.

$$\sup_{z \in \mathbb{C}} \|F(z)\|_X < \infty.$$

Then F is constant.

This is a generalization of Liouville’s theorem to the case where the target of the map is a Banach space. To prove this, we reduce this to the case of complex-valued functions. To do so, we compose this F with a map $X \rightarrow \mathbb{C}$.

Proof. Let $f \in X^*$. Then we show $f \circ F : \mathbb{C} \rightarrow \mathbb{C}$ is bounded and entire. To see it is bounded, just note that f is a bounded linear map. So

$$\sup_{z \in \mathbb{C}} |f \circ F(z)| \leq \sup_{z \in \mathbb{C}} \|f\|_{X^*} \|F(z)\|_X < \infty.$$

Analyticity can be shown in a similar fashion, exploiting the fact that f^* is linear.

Hence Liouville’s theorem implies $f \circ F$ is constant, i.e. $(f \circ F)(z) = (f \circ F)(0)$. In particular, this implies $f(F(z) - F(0)) = 0$. Moreover, this is true for all $f \in X^*$. Hence by (corollary of) Hahn-Banach theorem, we know $F(z) - F(0) = 0$ for all $z \in \mathbb{C}$. Therefore F is constant. \square

We have thus completed our proof that $\sigma(T)$ is non-empty, closed and a subset of $\{\lambda \in \mathbb{C} : |\lambda| \leq \|T\|_{\mathcal{B}(X)}\}$.

However, we want to know more. Apart from the spectrum itself, we also had the point spectrum $\sigma_p(T)$ and the approximate point spectrum $\sigma_{ap}(T)$, and we had the inclusions

$$\sigma_p(T) \subseteq \sigma_{ap}(T) \subseteq \sigma(T).$$

We know that the largest set $\sigma(T)$ is non-empty, but we want the smaller ones to be non-empty as well. We have the following theorem:

Theorem. We have

$$\sigma_{ap}(T) \supseteq \partial\sigma(T),$$

where $\partial\sigma(T)$ is the boundary of $\sigma(T)$ in the topology of \mathbb{C} . In particular, $\sigma_{ap}(T) \neq \emptyset$.

On the other hand, it is possible for $\sigma_p(T)$ to be empty (in infinite dimensional cases).

Proof. Let $\lambda \in \partial\sigma(T)$. Pick sequence $\{\lambda_n\}_{n=1}^\infty \subseteq \rho(T) = \mathbb{C} \setminus \sigma(T)$ such that $\lambda_n \rightarrow \lambda$. We claim that $R(\lambda_n) = (T - \lambda_n I)^{-1}$ satisfies

$$\|R(\lambda_n)\|_{\mathcal{B}(X)} \rightarrow \infty.$$

If this were the case, then we can pick $\mathbf{y}_n \in X$ such that $\|\mathbf{y}_n\| \rightarrow 0$ and $\|R(\lambda_n)(\mathbf{y}_n)\| = 1$. Setting $\mathbf{x}_n = R(\lambda_n)(\mathbf{y}_n)$, we have

$$\begin{aligned} \|(T - \lambda I)\mathbf{x}_n\| &\leq \|(T - \lambda_n I)\mathbf{x}_n\|_X + \|(\lambda - \lambda_n)\mathbf{x}_n\|_X \\ &= \|(T - \lambda_n I)(T - \lambda_n I)^{-1}\mathbf{y}_n\|_X + \|(\lambda - \lambda_n)\mathbf{x}_n\| \\ &= \|\mathbf{y}_n\|_X + |\lambda - \lambda_n| \\ &\rightarrow 0. \end{aligned}$$

So $\lambda \in \sigma_{ap}(T)$.

Thus, it remains to prove that $\|R(\lambda_n)\|_{\mathcal{B}(X)} \rightarrow \infty$. Recall from last time if S_1 is invertible, and

$$\|S_1^{-1}\|_{\mathcal{B}(X)} \|S_1 - S_2\|_{\mathcal{B}(X)} \leq 1, \quad (*)$$

then S_2 is invertible. Thus, for any $\mu \in \sigma(T)$, we have

$$\|R(\lambda_n)\|_{\mathcal{B}(X)} |\mu - \lambda_n| = \|R(\lambda_n)\|_{\mathcal{B}(X)} \|(T - \lambda_n I) - (T - \mu I)\|_{\mathcal{B}(X)} \geq 1.$$

Thus, it follows that

$$\|R(\lambda_n)\|_{\mathcal{B}(X)} \geq \frac{1}{\inf\{|\mu - \lambda_n| : \mu \in \sigma(T)\}} \rightarrow \infty.$$

So we are done. □

Having proven so many theorems, we now look at an specific example.

Example. Consider the shift operator $S : \ell_\infty \rightarrow \ell_\infty$ defined by

$$(a_1, a_2, a_3, \dots) \mapsto (0, a_1, a_2, \dots).$$

Then S is a bounded linear operator with norm $\|S\|_{\mathcal{B}(\ell_\infty)} = 1$. The theorem then tells $\sigma(S)$ is a non-empty closed subset of $\{\lambda \in \mathbb{C} : \|\lambda\| \leq 1\}$.

First, we want to understand what the point spectrum is. In fact, it is empty. To show this, suppose

$$S(a_1, a_2, a_3, \dots) = \lambda(a_1, a_2, a_3, \dots)$$

for some $\lambda \in \mathbb{C}$. In other words,

$$(0, a_1, a_2, \dots) = \lambda(a_1, a_2, a_3, \dots).$$

First consider the possibility that $\lambda = 0$. This would imply that the left is zero. So $a_i = 0$ for all i .

If $\lambda \neq 0$, then for the first coordinate to match, we must have $a_1 = 0$. Then for the second coordinate to match, we also need $a_2 = 0$. By induction, we need all $a_i = 0$. So $\ker(S - \lambda I) = \{0\}$ for all $\lambda \in \mathbb{C}$.

To find the spectrum, we will in fact show that

$$\sigma(S) = D = \{\lambda \in \mathbb{C} : |\lambda| \leq 1\}.$$

To prove this, we need to show that for any $\lambda \in D$, $S - \lambda I$ is not surjective. The $\lambda = 0$ case is obvious. For the other cases, we first have a look at what the image of $S - \lambda I$ looks like. We take

$$(b_1, b_2, b_3, \dots) \in \ell_\infty.$$

Suppose for some $\lambda \in D$, there exists (a_1, a_2, \dots) such that we have

$$(S - \lambda I)(a_1, a_2, \dots) = (b_1, b_2, \dots)$$

In other words, we have

$$(0, a_1, a_2, \dots) - (\lambda a_1, \lambda a_2, \lambda a_3, \dots) = (b_1, b_2, b_3, \dots).$$

So $-\lambda a_1 = b_1$. Hence we have

$$a_1 = -\lambda^{-1}b_1.$$

The next line then gives

$$a_1 - \lambda a_2 = b_2.$$

Hence

$$a_2 = -\lambda^{-1}(b_2 - a_1) = -\lambda^{-1}(b_2 + \lambda^{-1}b_1).$$

Inductively, we can show that

$$a_n = \lambda^{-1}(b_n + \lambda^{-1}b_{n-1} + \lambda^{-2}b_{n-2} + \dots + \lambda^{-n+1}b_1).$$

Now if $|\lambda| \leq 1$, we pick b_n such that $|b_n| = 1$ and $\lambda^{-i}b_{n-i} = |\lambda|^{-i}$. Then we must have $|a_n| \rightarrow \infty$. Such a sequence $(a_n) \notin \ell_\infty$. So $(b_n) \notin \text{im}(S - \lambda I)$. Therefore for $|\lambda| \leq 1$, $S - \lambda I$ is not surjective.

Hence we have $\sigma(S) \supseteq D$. By the theorem, we also know $\sigma(S) \subseteq D$. So in fact $\sigma(S) = D$.

Finally, we show that

$$\sigma_{ap}(S) = \partial D = \{\lambda \in \mathbb{C} : |\lambda| = 1\}.$$

Our theorem tells us that $\partial D \subseteq \sigma_{ap}(S) \subseteq D$. To show that indeed $\partial D = \sigma_{ap}(S)$, note that if $|\lambda| < 1$, then for all $\mathbf{x} \in \ell_\infty$.

$$\|(S - \lambda I)\mathbf{x}\|_{\ell_\infty} \geq \|S\mathbf{x}\|_{\ell_\infty} - |\lambda|\|\mathbf{x}\|_{\ell_\infty} = \|\mathbf{x}\|_{\ell_\infty} - |\lambda|\|\mathbf{x}\|_{\ell_\infty} = (1 - |\lambda|)\|\mathbf{x}\|_{\ell_\infty}.$$

So if $|\lambda| < 1$, then there exists no sequence \mathbf{x}_n with $\|\mathbf{x}_n\|_{\ell_\infty} = 1$ and $\|(S - \lambda I)\mathbf{x}_n\|_{\ell_\infty} \rightarrow 0$. So λ is not in the approximate point spectrum.

We see that these results are rather unpleasant and the spectrum behaves rather unlike the finite dimensional cases. We saw last time that the spectrum $\sigma(T)$ can in general be complicated. In fact, any non-empty compact subset of \mathbb{C} can be realized as the spectrum of some operator T on some Hilbert space H . This is an exercise on the Example sheet.

We are now going to introduce a class of “nice” operators whose spectrum behaves in a way more similar to the finite-dimensional case. In particular, most of $\sigma(T)$ consists of $\sigma_p(T)$ and is discrete. This includes, at least, all finite rank operators (i.e. operators T such that $\dim(\text{im } T) < \infty$) and their limits.

Definition (Compact operator). Let X, Y be Banach spaces. We say $T \in \mathcal{L}(X, Y)$ is *compact* if for every bounded subset E of X , $T(E)$ is totally bounded.

We write $\mathcal{B}_0(X)$ for the set of all compact operators $T \in \mathcal{B}(X)$.

Note that in the definition of T , we only required T to be a linear map, not a bounded linear map. However, boundedness comes from the definition for free because a totally bounded set is bounded.

There is a nice alternative characterization of compact operators:

Proposition. Let X, Y be Banach spaces. Then $T \in \mathcal{L}(X, Y)$ is compact if and only if $T(B(1))$ is totally bounded if and only if $\overline{T(B(1))}$ is compact.

The first equivalence is obvious, since $B(1)$ is a bounded set, and given any bounded set E , we can rescale it to be contained in $B(1)$. The second equivalence comes from the fact that a space is compact if and only if it is totally bounded and complete.

The last characterization is what we will use most of the time, and this is where the name “compact operator” came from.

Proposition. Let X be a Banach space. Then $\mathcal{B}_0(X)$ is a closed subspace of $\mathcal{B}(X)$. Moreover, if $T \in \mathcal{B}_0(X)$ and $S \in \mathcal{B}(X)$, then $TS, ST \in \mathcal{B}_0(X)$.

In a more algebraic language, this means $\mathcal{B}_0(X)$ is a closed ideal of the algebra $\mathcal{B}(X)$.

Proof. There are three things to prove. First, it is obvious that $\mathcal{B}_0(X)$ is a subspace. To check it is closed, suppose $\{T_n\}_{n=1}^\infty \subseteq \mathcal{B}_0(X)$ and $\|T_n - T\|_{\mathcal{B}(X)} \rightarrow 0$. We need to show $T \in \mathcal{B}_0(X)$, i.e. $T(B(1))$ is totally bounded.

Let $\varepsilon > 0$. Then there exists N such that

$$\|T - T_n\|_{\mathcal{B}(X)} < \varepsilon$$

whenever $n \geq N$. Take such an n . Then $T_n(B(1))$ is totally bounded. So there exists $\mathbf{x}_1, \dots, \mathbf{x}_k \in B(1)$ such that $\{T_n \mathbf{x}_i\}_{i=1}^k$ is an ε -net for $T_n(B(1))$. We now claim that $\{T \mathbf{x}_i\}_{i=1}^k$ is a 3ε -net for $T(B(1))$.

This is easy to show. Let $\mathbf{x} \in X$ be such that $\|\mathbf{x}\| \leq 1$. Then by the triangle inequality,

$$\begin{aligned} \|T\mathbf{x} - T\mathbf{x}_i\|_X &\leq \|T\mathbf{x} - T_n\mathbf{x}\| + \|T_n\mathbf{x} - T_n\mathbf{x}_i\| + \|T_n\mathbf{x}_i - T\mathbf{x}_i\| \\ &\leq \varepsilon + \|T_n\mathbf{x} - T_n\mathbf{x}_i\|_X + \varepsilon \\ &= 2\varepsilon + \|T_n\mathbf{x} - T_n\mathbf{x}_i\|_X \end{aligned}$$

Now since $\{T_n \mathbf{x}_i\}$ is an ε -net for $T_n(B(1))$, there is some i such that $\|T_n \mathbf{x} - T_n \mathbf{x}_i\| < \varepsilon$. So this gives

$$\|T\mathbf{x} - T\mathbf{x}_i\|_X \leq 3\varepsilon.$$

Finally, let $T \in \mathcal{B}_0(X)$ and $S \in \mathcal{B}(X)$. Let $\{\mathbf{x}_n\} \subseteq X$ such that $\|\mathbf{x}_n\|_X \leq 1$. Since T is compact, i.e. $\overline{T(B(1))}$ is compact, there exists a convergence subsequence of $\{T\mathbf{x}_i\}$.

Since S is bounded, it maps a convergent sequence to a convergent sequence. So $\{ST\mathbf{x}_n\}$ also has a convergent subsequence. So $\overline{ST(B(1))}$ is compact. So ST is compact.

We also have to show that $TS(B(1))$ is totally bounded. Since S is bounded, $S(B(1))$ is bounded. Since T sends a bounded set to a totally bounded set, it follows that $TS(B(1))$ is totally bounded. So TS is compact. \square

At this point, it helps to look at some examples at actual compact operators.

Example.

- (i) Let X, Y be Banach spaces. Then any finite rank operator in $\mathcal{B}(X, Y)$ is compact. This is since $T(B(1))$ is a bounded subset of a finite-dimensional space (since T is bounded), and any bounded subset of a finite-dimensional space is totally bounded.

In particular, any $f \in X^*$ is compact. Moreover, by the previous proposition, limits of finite-rank operators are also compact (since $\mathcal{B}_0(X)$ is closed).

- (ii) Let X be a Banach space. Then $I : X \rightarrow X$ is compact if and only if X is finite dimensional. This is since $\overline{B(1)}$ is compact if and only if the space is finite-dimensional.
- (iii) Let $K : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function. Define $T : C([0, 1]) \rightarrow C([0, 1])$ by the convolution

$$(Tf)(x) = \int_0^1 K(x-y)f(y) dy.$$

We first show T is bounded. This is since

$$\sup_x |Tf(x)| \leq \sup_{z \in [-1, 1]} K(z) \sup_y |f(y)|.$$

Since K is smooth, it is bounded on $[-1, 1]$. So Tf is bounded. In fact, T is compact. To see this, notice

$$\sup_x \left| \frac{d(Tf)}{dx}(x) \right| \leq \sup_{z \in [-1, 1]} |K'(z)| \sup_y |f(y)|.$$

Therefore if we have a sequence $\{f_n\} \subseteq C([0, 1])$ with $\|f_n\|_{C([0, 1])} \leq 1$, then $\{Tf_n\}_{n=1}^\infty$ is uniformly bounded with uniformly bounded derivative, and hence equicontinuous. By Arzelà-Ascoli theorem, $\{Tf_n\}_{n=1}^\infty$ has a convergent subsequence. Therefore $\overline{T(B(1))}$ is compact.

There is much more we can say about the last example. For example, requiring K to be smooth is clearly overkill. Requiring, say, differentiable with bounded derivative is already sufficient. Alternatively, we can ask what we can get if we work on, say, L^2 instead of $C([0, 1])$.

However, we don't have enough time for that, and instead we should return to developing some general theory. An important result is the following theorem, characterizing the point spectrum and spectrum of a compact operator.

Theorem. Let X be an infinite-dimensional Banach space, and $T \in \mathcal{B}(X)$ be a compact operator. Then $\sigma_p(T) = \{\lambda_i\}$ is at most countable. If $\sigma_p(T)$ is infinite, then $\lambda_i \rightarrow 0$.

The spectrum is given by $\sigma(T) = \sigma_p(T) \cup \{0\}$. Moreover, for every non-zero $\lambda_i \in \sigma_p(T)$, the eigenspace has finite dimensions.

Note that it is still possible for a compact operator to have empty point spectrum. In that case, the spectrum is just $\{0\}$. An example of this is found on the example sheet.

We will only prove this in the case where $X = H$ is a Hilbert space. In a lot of the proofs, we will have a closed subspace $V \leq X$, and we often want to pick an element in $X \setminus V$ that is “far away” from V in some sense. If we have a Hilbert space, then we can pick this element to be an element in the complement of V . If we are not in a Hilbert space, then we need to invoke *Riesz’s lemma*, which we shall not go into.

We will break the full proof of the theory into many pieces.

Proposition. Let H be a Hilbert space, and $T \in \mathcal{B}_0(H)$ a compact operator. Let $a > 0$. Then there are only finitely many linearly independent eigenvectors whose eigenvalue have magnitude $\geq a$.

This already gives most of the theorem, since this mandates $\sigma_p(T)$ is at most countable, and if $\sigma_p(T)$ is infinite, we must have $\lambda_i \rightarrow 0$. Since there are only finitely many linearly independent *eigenvectors*, the eigenspaces are also finite.

Proof. Suppose not. Then there are infinitely many independent $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$ such that $T\mathbf{x}_i = \lambda_i\mathbf{x}_i$ with $|\lambda_i| \geq a$.

Define $X_n = \text{span}\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$. Since the \mathbf{x}_i ’s are linearly independent, there exists $\mathbf{y}_n \in X_n \cap X_{n-1}^\perp$ with $\|\mathbf{y}_n\|_H = 1$.

Now let

$$\mathbf{z}_n = \frac{\mathbf{y}_n}{\lambda_n}.$$

Note that

$$\|\mathbf{z}_n\|_H \leq \frac{1}{a}.$$

Since X_n is spanned by the eigenvectors, we know that T maps X_n into itself. So we have

$$T\mathbf{z}_n \in X_n.$$

Moreover, we claim that $T\mathbf{z}_n - \mathbf{y}_n \in X_{n-1}$. We can check this directly. Let

$$\mathbf{y}_n = \sum_{k=1}^n c_k \mathbf{x}_k.$$

Then we have

$$\begin{aligned} T\mathbf{z}_n - \mathbf{y}_n &= \frac{1}{\lambda_n} T \left(\sum_{k=1}^n c_k \mathbf{x}_k \right) - \sum_{k=1}^n c_k \mathbf{x}_k \\ &= \sum_{k=1}^n c_k \left(\frac{\lambda_k}{\lambda_n} - 1 \right) \mathbf{x}_k \\ &= \sum_{k=1}^{n-1} c_k \left(\frac{\lambda_k}{\lambda_n} - 1 \right) \mathbf{x}_k \in X_{n-1}. \end{aligned}$$

We next claim that $\|T\mathbf{z}_n - T\mathbf{z}_m\|_H \geq 1$ whenever $n > m$. If this holds, then T is not compact, since $T\mathbf{z}_n$ does not have a convergent subsequence.

To show this, wlog, assume $n > m$. We have

$$\|T\mathbf{z}_n - T\mathbf{z}_m\|_H^2 = \|(T\mathbf{z}_n - \mathbf{y}_n) - (T\mathbf{z}_m - \mathbf{y}_n)\|_H^2$$

Note that $T\mathbf{z}_n - \mathbf{y}_n \in X_{n-1}$, and since $m < n$, we also have $T\mathbf{z}_m \in X_{n-1}$. By construction, $\mathbf{y}_n \perp X_{n-1}$. So by Pythagorean theorem, we have

$$\begin{aligned} &= \|T\mathbf{z}_n - \mathbf{y}_n - T\mathbf{z}_m\|_H^2 + \|\mathbf{y}_n\|_H^2 \\ &\geq \|\mathbf{y}_n\|_H^2 \\ &= 1 \end{aligned}$$

So done. □

To prove the previous theorem, the only remaining thing to prove is that $\sigma(T) = \sigma_p(T) \cup \{0\}$. In order to prove this, we need a lemma, which might seem a bit unmotivated at first, but will soon prove itself useful.

Lemma. Let H be a Hilbert space, and $T \in \mathcal{B}(H)$ compact. Then $\text{im}(I - T)$ is closed.

Proof. We let S be the orthogonal complement of $\ker(I - T)$, which is a closed subspace, hence a Hilbert space. We shall consider the restriction $(I - T)|_S$, which has the same image as $I - T$.

To show that $\text{im}(I - T)$ is closed, it suffices to show that $(I - T)|_S$ is bounded below, i.e. there is some $C > 0$ such that

$$\|\mathbf{x}\|_H \leq C\|(I - T)\mathbf{x}\|_H$$

for all $\mathbf{x} \in S$. If this were the case, then if $(I - T)\mathbf{x}_n \rightarrow \mathbf{y}$ in H , then

$$\|\mathbf{x}_n - \mathbf{x}_m\| \leq C\|(I - T)(\mathbf{x}_n - \mathbf{x}_m)\| \rightarrow 0,$$

and so $\{\mathbf{x}_n\}$ is a Cauchy sequence. Write $\mathbf{x}_n \rightarrow \mathbf{x}$. Then by continuity, $(I - T)\mathbf{x} = \mathbf{y}$, and so $\mathbf{y} \in \text{im}(I - T)$.

Thus, suppose $(I - T)$ is not bounded below. Pick \mathbf{x}_n such that $\|\mathbf{x}_n\|_H = 1$, but $(I - T)\mathbf{x}_n \rightarrow 0$. Since T is compact, we know $T\mathbf{x}_n$ has a convergent subsequence. We may wlog $T\mathbf{x}_n \rightarrow \mathbf{y}$. Then since $\|T\mathbf{x}_n - \mathbf{x}_n\|_H \rightarrow 0$, it follows that we also have $\mathbf{x}_n \rightarrow \mathbf{y}$. In particular, $\|\mathbf{y}\| = 1 \neq 0$, and $\mathbf{y} \in S$.

But $\mathbf{x}_n \rightarrow \mathbf{y}$ also implies $T\mathbf{x}_n \rightarrow T\mathbf{y}$. So this means we must have $T\mathbf{y} = \mathbf{y}$. But this is a contradiction, since \mathbf{y} does not lie in $\ker(I - T)$. □

Proposition. Let H be a Hilbert space, $T \in \mathcal{B}(H)$ compact. If $\lambda \neq 0$ and $\lambda \in \sigma(T)$, then $\lambda \in \sigma_p(T)$.

Proof. We will prove if $\lambda \neq 0$ and $\lambda \notin \sigma_p(T)$, then $\lambda \notin \sigma(T)$. In other words, let $\lambda \neq 0$ and $\ker(T - \lambda I) = \{0\}$. We will show that $T - \lambda I$ is surjective, i.e. $\text{im}(T - \lambda I) = H$.

Suppose this is not the case. Denote $H_0 = H$ and $H_1 = \text{im}(T - \lambda I)$. We know that H_1 is closed and is hence a Hilbert space. Moreover, $H_1 \subsetneq H_0$ by assumption.

We now define the sequence $\{H_n\}$ recursively by

$$H_n = (T - \lambda I)H_{n-1}.$$

We claim that $H_n \subsetneq H_{n-1}$. This must be the case, because the map $(T - \lambda I)^n : H_0 \rightarrow H_n$ is an isomorphism (it is injective and surjective). So the inclusion $H_n \subseteq H_{n-1}$ is isomorphic to the inclusion $H_1 \subseteq H_0$, which is strict.

Thus we have a strictly decreasing sequence

$$H_0 \supsetneq H_1 \supsetneq H_2 \supsetneq \cdots$$

Let \mathbf{y}_n be such that $\mathbf{y}_n \in H_n$, $\mathbf{y}_n \perp H_{n+1}$ and $\|\mathbf{y}_n\|_H = 1$. We now claim $\|T\mathbf{y}_n - T\mathbf{y}_m\| \geq |\lambda|$ if $n \neq m$. This then contradicts the compactness of T . To show this, again wlog we can assume that $n > m$. Then we have

$$\begin{aligned} \|T\mathbf{y}_n - T\mathbf{y}_m\|_H^2 &= \|(T\mathbf{y}_n - \lambda\mathbf{y}_n) - (T\mathbf{y}_m - \lambda\mathbf{y}_m) - \lambda\mathbf{y}_m + \lambda\mathbf{y}_n\|_H^2 \\ &= \|(T - \lambda I)\mathbf{y}_n - (T - \lambda I)\mathbf{y}_m - \lambda\mathbf{y}_m + \lambda\mathbf{y}_n\|_H^2 \end{aligned}$$

Now note that $(T - \lambda I)\mathbf{y}_n \in H_{n+1} \subseteq H_{m+1}$, while $(T - \lambda I)\mathbf{y}_m$ and $\lambda\mathbf{y}_n$ are both in H_{m+1} . So $\lambda\mathbf{y}_m$ is perpendicular to all of them, and Pythagorean theorem tells

$$\begin{aligned} &= |\lambda|^2 \|\mathbf{y}_m\|^2 + \|(T - \lambda I)\mathbf{y}_n - (T - \lambda I)\mathbf{y}_m - \lambda\mathbf{y}_m\|^2 \\ &\geq |\lambda|^2 \|\mathbf{y}_m\|^2 \\ &= |\lambda|^2. \end{aligned}$$

This contradicts the compactness of T . Therefore $\text{im}(T - \lambda I) = H$. \square

Finally, we can prove the initial theorem.

Theorem. Let H be an infinite-dimensional Hilbert space, and $T \in \mathcal{B}(H)$ be a compact operator. Then $\sigma_p(T) = \{\lambda_i\}$ is at most countable. If $\sigma_p(T)$ is infinite, then $\lambda_i \rightarrow 0$.

The spectrum is given by $\sigma(T) = \sigma_p(T) \cup \{0\}$. Moreover, for every non-zero $\lambda_i \in \sigma_p(T)$, the eigenspace has finite dimensions.

Proof. As mentioned, it remains to show that $\sigma(T) = \sigma_p(T) \cup \{0\}$. The previous proposition tells us $\sigma(T) \setminus \{0\} \subseteq \sigma_p(T)$. So it only remains to show that $0 \in \sigma(T)$.

There are two possible cases. The first is if $\{\lambda_i\}$ is infinite. We have already shown that $\lambda_i \rightarrow 0$. So $0 \in \sigma(T)$ by the closedness of the spectrum.

Otherwise, if $\{\lambda_i\}$ is finite, let $E_{\lambda_1}, \dots, E_{\lambda_n}$ be the eigenspaces. Define

$$H' = \text{span}\{E_{\lambda_1}, \dots, E_{\lambda_n}\}^\perp.$$

This is non-empty, since each E_{λ_i} is finite-dimensional, but H is infinite dimensional. Then T restricts to $T|_{H'} : H' \rightarrow H'$.

Now $T|_{H'}$ has no non-zero eigenvalues. By the previous discussion, we know $\sigma(T|_{H'}) \subseteq \{0\}$. By non-emptiness of $\sigma(T|_{H'})$, we know $0 \in \sigma(T|_{H'}) \subseteq \sigma(T)$.

So done. \square

4.6 Self-adjoint operators

We have just looked at compact operators. This time, we are going to add a condition of self-adjointness.

Definition (Self-adjoint operator). Let H be a Hilbert space, $T \in \mathcal{B}(H)$. Then T is *self-adjoint* or *Hermitian* if for all $\mathbf{x}, \mathbf{y} \in H$, we have

$$\langle T\mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{x}, T\mathbf{y} \rangle.$$

It is important to note that we defined the term for bounded linear operators T . If we have unbounded operators instead, Hermitian means something different from self-adjoint, and we have to be careful.

Recall that we defined the adjoint of a linear map to be a map of the dual spaces. However, we will often abuse notation and call $T^* : H \rightarrow H$ the *adjoint*, which is the (unique) operator such that for all $\mathbf{x}, \mathbf{y} \in H$,

$$\langle T\mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{x}, T^*\mathbf{y} \rangle.$$

It is an exercise to show that this is well-defined.

How is this related to the usual adjoint? Let $\tilde{T}^* : H^* \rightarrow H^*$ be the usual adjoint. Then we have

$$T^* = \phi^{-1} \circ \tilde{T}^* \circ \phi,$$

where $\phi : H \rightarrow H^*$ is defined by

$$\phi(\mathbf{v})(\mathbf{w}) = \langle \mathbf{w}, \mathbf{v} \rangle$$

as in the Riesz representation theorem.

The main result regarding self-adjoint operators is the spectral theorem:

Theorem (Spectral theorem). Let H be an infinite dimensional Hilbert space and $T : H \rightarrow H$ a compact self-adjoint operator.

- (i) $\sigma_p(T) = \{\lambda_i\}_{i=1}^N$ is at most countable.
- (ii) $\sigma_p(T) \subseteq \mathbb{R}$.
- (iii) $\sigma(T) = \{0\} \cup \sigma_p(T)$.
- (iv) If E_{λ_i} are the eigenspaces, then $\dim E_{\lambda_i}$ is finite if $\lambda_i \neq 0$.
- (v) $E_{\lambda_i} \perp E_{\lambda_j}$ if $\lambda_i \neq \lambda_j$.
- (vi) If $\{\lambda_i\}$ is infinite, then $\lambda_i \rightarrow 0$.
- (vii)

$$T = \sum_{i=1}^N \lambda_i P_{E_{\lambda_i}}.$$

We have already shown (i), (iii), (iv) and (vi). The parts (ii) and (v) we already did in IA Vectors and Matrices, but for completeness, we will do the proof again. They do not require compactness. The only non-trivial bit left is the last part (vii).

We first do the two easy bits.

Proposition. Let H be a Hilbert space and $T \in \mathcal{B}(H)$ self-adjoint. Then $\sigma_p(T) \subseteq \mathbb{R}$.

Proof. Let $\lambda \in \sigma_p(T)$ and $\mathbf{v} \in \ker(T - \lambda I) \setminus \{0\}$. Then by definition of \mathbf{v} , we have

$$\lambda = \frac{\langle T\mathbf{v}, \mathbf{v} \rangle}{\|\mathbf{v}\|_H^2} = \frac{\langle \mathbf{v}, T\mathbf{v} \rangle}{\|\mathbf{v}\|_H^2} = \bar{\lambda}.$$

So $\lambda \in \mathbb{R}$. □

Proposition. Let H be a Hilbert space and $T \in \mathcal{B}(H)$ self-adjoint. If $\lambda, \mu \in \sigma_p(T)$ and $\lambda \neq \mu$, then $E_\lambda \perp E_\mu$.

Proof. Let $\mathbf{v} \in \ker(T - \lambda I) \setminus \{0\}$ and $\mathbf{w} \in \ker(T - \mu I) \setminus \{0\}$. Then

$$\lambda \langle \mathbf{v}, \mathbf{w} \rangle = \langle T\mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{v}, T\mathbf{w} \rangle = \bar{\mu} \langle \mathbf{v}, \mathbf{w} \rangle = \mu \langle \mathbf{v}, \mathbf{w} \rangle,$$

using the fact that eigenvalues are real. Since $\lambda \neq \mu$ by assumption, we must have $\langle \mathbf{v}, \mathbf{w} \rangle = 0$. □

To prove the final part, we need the following proposition:

Proposition. Let H be a Hilbert space and $T \in \mathcal{B}(H)$ a compact self-adjoint operator. If $T \neq 0$, then T has a non-zero eigenvalue.

This is consistent with our spectral theorem, since if T is non-zero, then something in the sum $\lambda_i P_{E_{\lambda_i}}$ has to be non-zero. It turns out this is most of the work we need.

However, to prove this, we need the following lemma:

Lemma. Let H be a Hilbert space, and $T \in \mathcal{B}(H)$ a compact self-adjoint operator. Then

$$\|T\|_{\mathcal{B}(H)} = \sup_{\|\mathbf{x}\|_H=1} |\langle \mathbf{x}, T\mathbf{x} \rangle|$$

Proof. Write

$$\lambda = \sup_{\|\mathbf{x}\|_H=1} |\langle \mathbf{x}, T\mathbf{x} \rangle|.$$

Note that one direction is easy, since for all \mathbf{x} , Cauchy-Schwarz gives

$$|\langle \mathbf{x}, T\mathbf{x} \rangle| \leq \|T\mathbf{x}\|_H \|\mathbf{x}\|_H = \|T\|_{\mathcal{B}(H)} \|\mathbf{x}\|_H^2.$$

So it suffices to show the inequality in the other direction. We now claim that

$$\|T\|_{\mathcal{B}(H)} = \sup_{\|\mathbf{x}\|_H=1, \|\mathbf{y}\|_H=1} |\langle T\mathbf{x}, \mathbf{y} \rangle|.$$

To show this, recall that $\phi : H \rightarrow H^*$ defined by $\mathbf{v} \mapsto \langle \cdot, \mathbf{v} \rangle$ is an isometry. By definition, we have

$$\|T\|_{\mathcal{B}(H)} = \sup_{\|\mathbf{x}\|_H=1} \|T\mathbf{x}\|_H = \sup_{\|\mathbf{x}\|_H=1} \|\phi(T\mathbf{x})\|_{H^*} = \sup_{\|\mathbf{x}\|_H=1} \sup_{\|\mathbf{y}\|_H=1} |\langle \mathbf{y}, T\mathbf{x} \rangle|.$$

Hence, it suffices to show that

$$\sup_{\|\mathbf{x}\|_H=1, \|\mathbf{y}\|_H=1} |\langle T\mathbf{x}, \mathbf{y} \rangle| \leq \lambda.$$

Take $\mathbf{x}, \mathbf{y} \in H$ such that $\|\mathbf{x}\|_H = \|\mathbf{y}\|_H = 1$. We first perform a trick similar to the polarization identity. First, by multiplying \mathbf{y} by an appropriate scalar, we can wlog assume $\langle T\mathbf{x}, \mathbf{y} \rangle$ is real. Then we have

$$\begin{aligned} |\langle T(\mathbf{x} + \mathbf{y}), \mathbf{x} + \mathbf{y} \rangle - \langle T(\mathbf{x} - \mathbf{y}), \mathbf{x} - \mathbf{y} \rangle| &= 2|\langle T\mathbf{x}, \mathbf{y} \rangle + \langle T\mathbf{y}, \mathbf{x} \rangle| \\ &= 4|\langle T\mathbf{x}, \mathbf{y} \rangle|. \end{aligned}$$

Hence we have

$$\begin{aligned} |\langle T\mathbf{x}, \mathbf{y} \rangle| &= \frac{1}{4} |\langle T(\mathbf{x} + \mathbf{y}), \mathbf{x} + \mathbf{y} \rangle - \langle T(\mathbf{x} - \mathbf{y}), \mathbf{x} - \mathbf{y} \rangle| \\ &\leq \frac{1}{4} (\lambda \|\mathbf{x} + \mathbf{y}\|_H^2 + \lambda \|\mathbf{x} - \mathbf{y}\|_H^2) \\ &= \frac{\lambda}{4} (2\|\mathbf{x}\|_H^2 + 2\|\mathbf{y}\|_H^2) \\ &= \lambda, \end{aligned}$$

where we used the parallelogram law. So we have $\|T\|_{\mathcal{B}(H)} \leq \lambda$. \square

Finally, we can prove our proposition.

Proposition. Let H be a Hilbert space and $T \in \mathcal{B}(H)$ a compact self-adjoint operator. If $T \neq 0$, then T has a non-zero eigenvalue.

Proof. Since $T \neq 0$, then $\|T\|_{\mathcal{B}(H)} \neq 0$. Let $\|T\|_{\mathcal{B}(H)} = \lambda$. We now claim that either λ or $-\lambda$ is an eigenvalue of T .

By the previous lemma, there exists a sequence $\{\mathbf{x}_n\}_{n=1}^\infty \subseteq H$ such that $\|\mathbf{x}_n\|_H = 1$ and $\langle \mathbf{x}_n, T\mathbf{x}_n \rangle \rightarrow \pm\lambda$.

We consider the two cases separately. Suppose $\langle \mathbf{x}_n, T\mathbf{x}_n \rangle \rightarrow \lambda$. Consider $T\mathbf{x}_n - \lambda\mathbf{x}_n$. Since T is compact, there exists a subsequence such that $T\mathbf{x}_{n_k} \rightarrow \mathbf{y}$ for some $\mathbf{y} \in H$. For simplicity of notation, we assume $T\mathbf{x}_n \rightarrow \mathbf{y}$ itself. We have

$$\begin{aligned} 0 &\leq \|T\mathbf{x}_n - \lambda\mathbf{x}_n\|_H^2 \\ &= \langle T\mathbf{x}_n - \lambda\mathbf{x}_n, T\mathbf{x}_n - \lambda\mathbf{x}_n \rangle \\ &= \|T\mathbf{x}_n\|_H^2 - 2\lambda\langle T\mathbf{x}_n, \mathbf{x}_n \rangle + \lambda^2\|\mathbf{x}_n\|^2 \\ &\rightarrow \lambda^2 - 2\lambda^2 + \lambda^2 \\ &= 0 \end{aligned}$$

as $n \rightarrow \infty$. Note that we implicitly used the fact that $\langle T\mathbf{x}_n, \mathbf{x}_n \rangle = \langle \mathbf{x}_n, T\mathbf{x}_n \rangle$ since $\langle T\mathbf{x}_n, \mathbf{x}_n \rangle$ is real. So we must have

$$\|T\mathbf{x}_n - \lambda\mathbf{x}_n\|_H^2 \rightarrow 0.$$

In other words,

$$\mathbf{x}_n \rightarrow \frac{1}{\lambda}\mathbf{y}.$$

Finally, we show \mathbf{y} is an eigenvector. This is easy, since

$$T\mathbf{y} = \lim_{n \rightarrow \infty} T(\lambda\mathbf{x}_n) = \lambda\mathbf{y}.$$

The case where $\mathbf{x}_n \rightarrow -\lambda$ is entirely analogous. In this case, $-\lambda$ is an eigenvalue. The proof is exactly the same, apart from some switching of signs. \square

Finally, we can prove the last part of the spectral theorem.

Proposition. Let H be an infinite dimensional Hilbert space and $T : H \rightarrow H$ a compact self-adjoint operator. Then

$$T = \sum_{i=1}^N \lambda_i P_{E_{\lambda_i}}.$$

Proof. Let

$$U = \text{span}\{E_{\lambda_1}, E_{\lambda_2}, \dots\}.$$

Firstly, we clearly have

$$T|_U = \sum_{i=1}^N \lambda_i P_{E_{\lambda_i}}.$$

This is since for any $\mathbf{x} \in U$ can be written as

$$\mathbf{x} = \sum_{i=1}^n P_{E_{\lambda_i}} \mathbf{x}.$$

Less trivially, this is also true for \bar{U} , i.e.

$$T|_{\bar{U}} = \sum_{i=1}^N \lambda_i P_{E_{\lambda_i}},$$

but this is also clear from definition once we stare at it hard enough.

We also know that

$$H = \bar{U} \oplus U^\perp.$$

It thus suffices to show that

$$T|_{U^\perp} = 0.$$

But since $T|_{U^\perp}$ has no non-zero eigenvalues, this follows from our previous proposition. So done. \square